arXiv:1105.5471v1 [math.FA] 27 May 2011

ALGEBRAS OF SEMICLASSICAL PSEUDODIFFERENTIAL OPERATORS ASSOCIATED WITH ZOLL-TYPE DOMAINS IN COTANGENT BUNDLES

GERARDO HERNÁNDEZ-DUEÑAS* AND ALEJANDRO URIBE**

ABSTRACT. We are consider domains in cotangent bundles with the property that the null foliation of their boundary is fibrating and the leaves satisfy a Bohr-Sommerfeld condition (for example, the unit disk bundle of a Zoll metric). Given such a domain, we construct an algebra of associated semiclassical pseudodifferential operators with singular symbols. The Schwartz kernels of the operators have frequency set contained in the union of the diagonal and the flow-out of the null foliation of the boundary of the domain. We develop a symbolic calculus, prove the existence of projectors (under a mild additional assumption) whose range can be thought of as quantizing the domain, give a symbolic proof of a Szegö limit theorem, and study associated propagators.

CONTENTS

1. Introduction 2
1.1. Main results 4
2. The model case 6
2.1. Definitions 6
2.2. The symbol maps 8
3. The manifold case 12
3.1. The spaces $J^{\ell,m}$ on manifolds 12
3.2. Symbolic calculus 15
3.3. The existence of the symbolic calculus. 16
3.4. A symbolic compatibility condition 18
3.5. The adjoint 21
4. Asymptotics of the trace 22
4.1. The trace in case $m' \geq 0$. 22
4.2. The trace in case $m' \leq -4$. 28
5. Projectors and "cut" quantum observables 28
5.1. On the Existence of Projectors 29
5.2. Cut quantum observables 32

---

*Work supported in part by CONACYT #160147.
**Work supported in part by NSF grant DMS-0805878.

6. On some propagators $e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}$ 38
6.1. The classical counterpart 38
6.2. A Symbolic Description of The Propagator $e^{-it\ h^{-1}\Pi\widehat{Q}\Pi}$ 41
6.3. An Egorov-type Theorem 45
7. A numerical study of propagation of coherent states 46
References 48

1. INTRODUCTION

Let M be a smooth compact manifold, and $X_c \subset X := T^*M$ a closed compact domain with a smooth boundary. In this paper we address the question: what quantum objects are naturally associated with X_c? This question has been addressed indirectly in "classical" cases. For example, if X_c is the unit disk bundle associated to a Riemannian metric on M assumed compact, then morally speaking to X_c corresponds the ($\hbar$-dependent) subspace $\mathcal{H}$ of $L^2(M)$ spanned by the eigenfunctions of the semiclassical Laplacian, $\widehat{P} = \hbar^2\Delta$, with eigenvalues in $[0,1]$. This heuristics is in good measure justified by the theorem that, if $\Pi : L^2(M) \to \mathcal{H}$ is the orthogonal projection, then for any $\hbar$-pseudodifferential operator $\widehat{Q}$ one has:

$$\lim_{\hbar\to 0} \frac{1}{\dim \mathcal{H}}\ \operatorname{Tr} \Pi\widehat{Q}\Pi = \frac{1}{\mathrm{Vol} X_c} \int_{X_c} Q\, d\lambda, \tag{1.1}$$

where Q is the principal symbol of $\widehat{Q}$ and $d\lambda$ is Liouville measure (see [12]). But can one say more?

We discuss a more systematic answer to this question in the case when X_c is "fibrating and Bohr- Sommerfeld", by which we mean the following. The boundary ∂X_c is always foliated by curves tangent to the kernel of the pull-back of the symplectic form. The fibrating condition is that there exists a manifold S and a submersion $\pi : \partial X_c \to S$ whose fibers are the leaves of the null-foliation. This is satisfied iff X_c has a globally defining function whose Hamilton flow on ∂X_c is periodic with a common minimal period. The Bohr-Sommerfeld condition is that all leaves, γ, of π satisfy

$$\int_\gamma \alpha \in 2\pi\mathbb{Z},$$

where α is the tautological one-form in T^*M. We will show that under these conditions there exist spaces $J^{\ell,m}$ of pseudodifferential operators with singular symbols naturally associated with X_c. (Here (ℓ, m) is a bi-degree, to be explained later). The frequency sets of their Schwartz kernels are contained in the union of Lagrangian submanifolds of $T^*(M \times M)$

$$\Delta'_c \cup \mathcal{F}\partial X'_c,$$

where

$$\Delta_c = \{(\overline{x}, \overline{x}) \in X_c \times X_c\}$$

and $\mathcal{F}\partial X_c$ is the flow-out

$$\mathcal{F}\partial X_c = \{(\overline{x}, \overline{y}) \in \partial X_c \times \partial X_c \;;\; \overline{x},\ \overline{y} \text{ in the same leaf } \}.$$

Intuitively speaking, the diagonal part, Δ_c, is expected to be a part of any pseudodifferential operator calculus associated with X_c. The flow-out part, $\mathcal{F}\partial X_c$, is there because the fibrating-and-Bohr-Sommerfeld conditions imply that there should be a significant part of Hilbert space associated with the symplectic reduction of the boundary of X_c. The two symbols, one on Δ_c and one on $\mathcal{F}\partial X_c$, have a compatibility condition that comes about most naturally in our setting.

In particular, the space $J^{-1/2,1/2}$ is closed under composition, and it includes the projector Π mentioned above, in the case of Zoll metrics. The Bohr-Sommerfeld condition is needed for the existence of a global symbolic calculus, and goes along with having to restricting Planck's constant to take the values $\hbar = 1/N$, $N = 1, 2, \ldots$. The fibrating condition is needed in order for $\mathcal{F}\partial X_c$ to be a closed submanifold of $X \times X$.

The Schwartz kernels are semiclassical analogues of the oscillatory integrals with singular symbols of [22] and [11], associated to a pair of intersecting conic Lagrangian submanifolds. See also [19], where a precise calculus for a more generalized class is discussed, and [16] where a connection with a class of Legendre distributions is explained. In the conic case, the realization that if the Lagrangians are the diagonal and a flow-out one obtains a symbol calculus is due to the results in [1]. This is possible because

$$\Delta_c \circ \mathcal{F}\partial X_c,\ \mathcal{F}\partial X_c \circ \Delta_c,\ \mathcal{F}\partial X_c \circ \mathcal{F}\partial X_c \subset \mathcal{F}\partial X_c \;; \quad \Delta_c \circ \Delta_c \subset \Delta_c,$$

so composing two operators with wave-front set in $\Delta_c \cup \mathcal{F}\partial X_c$ produces an operator with wave-front set contained in the same union. We believe that the present semiclassical setting provides a very natural expression for the Antoniano-Uhlmann algebra.

In the Zoll case, there have been numerous papers aimed at refining the general result (1.1), mostly in terms of the reminder estimate. For instance, in [8], the second term in the Szegö formula for Zoll manifolds is proved. Other references in this direction are [9, 5, 21, 20, 24].

Here we are not focusing on remainder estimates, but on the fact that there is an operator algebra with a symbolic calculus that provides a quantization of X_c and, among other things, a broader (symbolic) setting for Szegö limit theorems. Furthermore, the existence of the operator algebra allows us to go farther in the analysis of the operators $\Pi\widehat{Q}\Pi$ mentioned above, with respect to previous works.

The issues we raise here are also connected with work of [7], on the relationship between symplectic cutting and quantization, in the homogeneous (non-semiclassical) category. They consider the case when $X_c = \phi^{-1}(-\infty, 0]$ where $\phi : X \to \mathbb{R}$ is the moment map for a homogeneous action of the circle group on S^1. In this setting one can form the symplectic cut

$$Y = X_c / \sim,$$

Their work centers on the algebra of operators $\{\Pi \widehat{Q} \Pi \ ; \ [\Pi, \widehat{Q}] = 0\}$ where $\widehat{Q}$ ranges over (non-semiclassical) ΨDOs on M and Π is a spectral projector, as above, associated to a quantization of the circle action by a Fourier integral operator. Roughly speaking they show that such an algebra can be considered a quantization of the symplectic manifold Y.

1.1. **Main results.** We now describe our main results.

Theorem 1.1. *Let $X_c \subset T^*M$ be as above (∂X_c "fibrating and Bohr-Sommerfeld"). Then there exist vector spaces of semiclassical pseudodifferential operators with singular symbols, $J^{\ell,m}(M \times M; \Delta, \mathcal{F}\partial X_c)$, where $\hbar$ is restricted to the sequence $1/N, N = 1, 2, \ldots$, such that:*

1. *The frequency set of the Schwartz kernel of any operator in the algebra is contained in the union $\Delta' \cup \mathcal{F}\partial X_c'$.*
2. *Let $\Sigma = \Delta \cap \mathcal{F}\partial X_c$. Then $J^{\ell,m}$ are microlocally Lagrangian states of order $\ell + m$ on $\Delta' \setminus \Sigma'$ and ℓ on $\mathcal{F}\partial X_c' \setminus \Sigma'$, and therefore there are symbol maps:*

   $$\sigma_0 : J \to |\wedge|^{1/2}(\Delta \setminus \Sigma), \quad \sigma_1 : J \to |\wedge|^{1/2}(\mathcal{F}\partial X_c \setminus \Sigma)$$

   (where $|\wedge|^{1/2}$ denotes the space of half-densities, and we will ignore Maslov factors). There is in fact a symbolic calculus, that will be described below.
3. *$J^{\ell,m} \circ J^{\tilde{\ell},\tilde{m}} \subset J^{\ell+\tilde{\ell}+1/2, m+\tilde{m}-1/2}$, in particular $J^{-1/2,1/2}$ is an algebra.*
4. *Assume that X_c is compact. Then every*

   $$\widehat{A} \in J^{-1/2,1/2}(M \times M; \Delta, \mathcal{F}\partial X_c)$$

   with microsupport contained in X_c is smoothing and

   $$\mathrm{Tr}(\widehat{A}) = (2\pi)^{-n} \ \hbar^{-n} \int_{X_c} \sigma_0(\widehat{A}) \, \frac{\omega^n}{n!} + O(\hbar^{-n+1} \log(1/\hbar))$$

   where ω is the symplectic form of T^*M and n is the dimension of M.

The manifold X_c has an associated operator algebra, $\mathcal{A}_{X_c}$, which consists of elements in the algebra $J\,(M \times M; \Delta, \partial X_c)$ which are microlocally of order $\mathrm{O}(\hbar^\infty)$ in the complement $T^*M \setminus X_c$.

Theorem 1.2. *(See §5.1 for details.) Assume ∂X_c is compact and of contact type. Then there exist orthogonal projections $\Pi \in J^{-1/2,1/2}$ whose symbol σ_0 is the characteristic function of X_c. Moreover, for any zeroth-order pseudodifferential operator on M, $\widehat{Q}$, the "cut" operator $\Pi\widehat{Q}\Pi$ is in the algebra, and σ_0 can be identified with $Q|_{X_c}$.*

As an immediate corollary we obtain the following Szegö limit theorem:

Corollary 1.3. *Assume that X_c is compact, $\Pi \in J^{-1/2,1/2}$ an orthogonal projector as in the previous theorem and let $\hbar = 1/N$. Then for any integer $m \geq 0$*

$$\mathrm{Tr}(\Pi\widehat{Q}\Pi)^m = (2\pi)^{-n}\ N^n \int_{X_c} Q^m \frac{\omega^n}{n!} + O(N^{n-1}\log(N))$$

Finally, we have a result on the propagator $\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}$ where the symbol, Q, of the pseudodifferential operator $\widehat{Q}$ preserves X_c "to second order":

Theorem 1.4. *Suppose $\widehat{Q}$ is a zeroth-order semiclassical pseudodifferential operator satisfying the conditions of Lemma 6.1. Assume sub$\widehat{Q}(\hbar) = 0$. Then*

$$\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi} \in J^{-1/2,1/2}\left(M \times M; \Delta(t), \mathcal{F}\partial X_c(t)\right),$$

where

$$\Delta(t) = \left\{(\overline{x},\overline{y})\,\middle|\,\overline{x},\overline{y} \in T^*(M \times M), \overline{x} = \Phi_t^Q(\overline{y})\right\}$$

$$\mathcal{F}\partial X_c(t) = \left\{(\overline{x},\overline{y})\,\middle|\,\overline{x},\overline{y} \in \partial X_c,\ \exists s \in \mathbb{R} \text{ such that } \overline{x} = \Phi_s^P\Phi_t^Q(\overline{y})\right\}.$$

Remarks:

1. The result on the trace (part (4) of the Theorem 1.1) extends to the spaces $J^{\ell,m}$ provided $m \geq -1/2$. If m is sufficiently negative, the leading contribution to the trace comes from the asymptotic singularity of the kernel of the operator at Σ.
2. The symbol calculus for σ_0 is just the usual pseudodifferential calculus. The symbol calculus for σ_1 is more complicated (and non-commutative). In fact σ_1 comes with an extension to a distribution on $\mathcal{F}\partial X_c$ conormal to Σ, and there exists a formula for the smooth part of the σ_1 of the composition in terms of the corresponding extensions of the factors.
3. The restriction that $\hbar = 1/N$ is necessary to have a well-defined global symbol on the flow-out $\mathcal{F}\partial X_c$, see [14] chapter VII, §0. Locally elements of the $J^{\ell,m}$ are given by semiclassical oscillatory integrals with $\hbar \to 0$ continuously.
4. The error estimate $O(\hbar\log(1/\hbar))$ is sharp for the class $J^{-1/2,1/2}$, but in Corollary 1.3 the error should be $O(\hbar^2)$, see Remark 4.4.

(5) The idea that the image of Π quantizes X_c and that the operators $\Pi\widehat{Q}\Pi$ can be considered as associated observables appeared first in the physics literature, see [2] §II E and [3] (where a connection with symplectic cutting is also made). The operators $\Pi\widehat{Q}\Pi$ do not form an algebra, however, while the class $J^{-1/2,1/2}$ does.

(6) Our work also implies that the results of [7] also hold in the semiclassical case.

The paper is organized as follows. In Section 2, the spaces $J^{\ell,m}$ are defined in a model case by oscillatory integrals with amplitudes having expansions in $\hbar$ with coefficients that are classical symbols. In Section 3, the spaces are defined globally on manifolds, and the existence of a symbolic calculus is established. The principal symbol on each Lagrangian blows-up at the intersection, and they satisfy a *symbolic* compatibility condition there. The theorem on the trace is proved in Section 4. Section 5 considers cases where the algebra admits projectors, yielding a symbolic proof of a generalized Szegö limit Theorem. Section 6 studies the propagator of certain elements in the algebra, and proves an Egorov-type theorem. Finally, Section 7 shows numerically a surprising phenomenon of propagation of coherent states in situations not considered in Section 6.

2. THE MODEL CASE

2.1. **Definitions.** We begin by discussing the microlocal model case: $M = \mathbb{R}^n$ and $X^n_c = \{p_1 \geq 0\}$, where $(x_1, \dots, x_n, p_1, \dots, p_n)$ are canonical coordinates in $T^*\mathbb{R}^n$. Let $\mathcal{F}\partial X^n_c$ be the flow-out of $\partial X^n_c = \{p_1 = 0\}$. We will use this case to define operators $J^{\ell,m}(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X^n_c)$, where $\Delta^n \subset T^*\mathbb{R}^n \times T^*\mathbb{R}^n$ is the diagonal.

Roughly speaking, elements in $J^{\ell,m}(\mathbb{R}^n \times \mathbb{R}^n, \Delta^n, \mathcal{F}\partial X^n_c)$ will be defined as oscillatory integrals with amplitudes as follows:

(1) We denote by $\mathcal{A}^{\ell,m}$ the class of all smooth functions $a(s, x, y, p, \sigma, \hbar)$ with compact support in s, x, y, p such that, as $\hbar \to 0$

$$a(s,x,y,p,\sigma,\hbar) \sim \sum_{j=-\ell}^{\infty} \hbar^j a_j(s,x,y,p,\sigma)$$

(in a sense that will be explained below) where, for each j, $a_j(s,x,y,p,\sigma)$ is a polyhomogeneous classical symbol in σ of degree m:

$$a_j(s,x,y,p,\sigma) \sim \sum_{r=m}^{-\infty} a_{j,r}(s,x,y,p,\sigma), \quad \forall\lambda > 0 \; a_{j,r}(s,x,y,p,\lambda\sigma) = \lambda^r a_{j,r}(s,x,y,p,\sigma).$$

(2) The operators in $J^{\ell,m}(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X^n_c)$ are those whose Schwartz kernels are of the form

(2.1)
$$A(x,y,\hbar) = \frac{1}{(2\pi\hbar)^n} \int e^{\frac{i}{\hbar}\left[(x_1-y_1-s)p_1 + (x'-y')p'\right] + is\sigma} \; a(s,x,y,p,\sigma,\hbar)\; ds\, dp\, d\sigma,$$

where we have split the variables: $x = (x_1, x'),\ y = (y_1, y')\ (x', y' \in \mathbb{R}^{n-1})$.

We now give the details.

Definition 2.1. Let (x, y) and s be the standard coordinates on $\mathbb{R}^{2n}$ and $\mathbb{R}$ respectively, and let p, σ be the dual coordinates to x, s. We denote by $z = (x, y, s)$ coordinates in $\mathbb{R}^{2n} \times \mathbb{R}$. Define $\mathcal{A}^{\ell,m}$ to be the space of smooth families $a(s, x, y, p, \sigma, \hbar)$ compactly supported in z, p such that

$$\left|(\partial/\partial z)^{\alpha} (\partial/\partial p)^{\beta} (\partial/\partial \sigma)^{\gamma} a\right| \leq C\hbar^{-\ell}(1+|\sigma|)^{m-|\gamma|}, \tag{2.2}$$

for some constant $C = C(\alpha, \beta, \gamma)$ and $\hbar \in (0, h_0]$ for some fixed $h_0 > 0$.

Let $a_j(s, x, y, p, \sigma) \in S^m$ be a sequence of classical symbols in the σ variable, of order m, and compactly supported in (s, x, y, p), i.e., a_j is smooth,

$$\left|(\partial/\partial z)^{\alpha} (\partial/\partial p)^{\beta} (\partial/\partial \sigma)^{\gamma} a_j(s, x, y, p, \sigma)\right| \leq C(\alpha, \beta, \gamma)\,(1+|\sigma|)^{m-|\gamma|},$$

for some constant $C = C(\alpha, \beta, \gamma)$, and there exists a sequence $a_{j,r}$ of smooth functions in $\sigma \neq 0$, homogeneous of degree r in σ, such that $a_j \sim \sum_{r=m}^{-\infty} a_{j,r}$ in the standard sense. Given $a \in \mathcal{A}^{\ell,m}$, we will say that $a \sim \sum_{j=-\ell}^{\infty} \hbar^j a_j$ iff for all integers $K \geq 0$

$$a - \sum_{j=-\ell}^{-\ell+K} \hbar^j a_j \in \mathcal{A}^{\ell-K-1,m}.$$

Definition 2.2. Denote by $\mathcal{A}^{\ell,m}_{\text{classical}} \subset \mathcal{A}^{\ell,m}$ the set of all $a(s, x, y, p, \sigma, \hbar)$ that satisfy $a \sim \sum_{j=-\ell}^{\infty} \hbar^j a_j$, as above, and let $\Sigma^n = \Delta^n \cap \mathcal{F}\partial X^n_c$. Define $J^{\ell,m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X^n_c\right)$ to be the set of kernels of the form

$$A(x, y, \hbar) + F_1(x, y, \hbar) + F_2(x, y, \hbar)$$

where $A(x, y, \hbar)$ is as in (2.1) with $a \in \mathcal{A}^{\ell',m'}_{\text{classical}}$ where

$$\ell' = \ell + 1/2, \quad m' = m - 1/2,$$

and F_1, F_2 are the kernels of semiclassical Fourier integral operator in $sc - I^{\ell+m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n \setminus \Sigma^n\right)$, $sc - I^{\ell}\left(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{F}\partial X^n_c\right)$, respectively.

Remark 2.3. As we will see, it is necessary that F_1 and F_2 appear in the definition in order for the classes J to be closed under composition.

We have yet to give sense to the formula in equation (2.1). Denote by $D = 1 + D_s^2$ where $D_s = \frac{1}{i}\frac{\partial}{\partial s}$. Notice that $De^{is\sigma} = (1+\sigma^2)e^{is\sigma}$. For $a \in A^{\ell',m'}_{\text{classical}}$, we define $A(x, y, \hbar)$ as

$$A(x, y, \hbar) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}(x-y)p + is\sigma} \frac{1}{(1+\sigma^2)^k} D^k \left(e^{-i\hbar^{-1}sp_1} a(s, x, y, p, \sigma, \hbar)\right) ds dp d\sigma. \tag{2.3}$$

The integral above is absolutely convergent for $k >> 0$ large enough since $D^k\left(e^{-i\hbar^{-1}sp_1}a\right)$ is $O(\sigma^m)$. It can be easily checked that the definition above does not depend on k using integration by parts.

Equivalently, one can define the integral in (2.1) as an iterated integral, where one first integrates over the variables (s,p) with respect to which the amplitude is compactly supported. The resulting function of σ is rapidly decreasing and therefore integrable. Both of these interpretations are useful in proofs.

We now state the first property of the kernels we have just defined:

Proposition 2.4. *For any $A \in J^{\ell,m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X_c^n\right)$, the frequency set of A is contained in the union of the Lagrangians Δ^n and $\mathcal{F}\partial X_c^n$:*

$$FS\,(A) \subset \Delta^{n\prime} \cup \mathcal{F}\partial X_c^{n\prime}.$$

Moreover, away from the intersection $\Sigma = \Delta^n \cap \mathcal{F}\partial X_c^n$, A is microlocally in the space sc-$I^{\ell+m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n \setminus \Sigma^n\right)$ and sc-$I^{\ell}\left(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{F}\partial X_c^n \setminus \Sigma^n\right)$, where sc-$I$ denote the spaces of semiclassical Fourier integral operators.

Remark 2.5. This is a consequence of Theorem 3.13, proved in Section 3.3.

2.2. **The symbol maps.** By the second part of Propositon 2.4, one has two symbol maps:

$$\begin{tikzcd} & {|\wedge|^{1/2}(\Delta^n \setminus \Sigma)} \\ J^{\ell,m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X_c^n\right) \arrow[ur, "\sigma_0"] \arrow[dr, "\sigma_1"'] & \\ & {|\wedge|^{1/2}(\mathcal{F}\partial X_c^n \setminus \Sigma).} \end{tikzcd} \tag{2.4}$$

It is easy to see that, for A as above, they are given by the following formulae:

$$\begin{aligned} &\sigma_0(A) := 2\pi\ a_{-\ell',m'}(s,x,y,p,\sigma)\sqrt{dxdp}\Big|_{y=x,s=0,p_1=\sigma} \qquad \text{and} \\ &\sigma_1(A) := \sqrt{2\pi}\ \int a_{-\ell'}(s,x,y,p,\sigma)e^{is\sigma}d\sigma\sqrt{dxdy_1dp'}\Big|_{y'=x',p_1=0,s=x_1-y_1}. \end{aligned} \tag{2.5}$$

Notice that σ_0 has a singularity as $\sigma = p_1$ converges to zero, that is, as the point where σ_0 is evaluated tends to the intersection, Σ. The same is true of σ_1, and the leading singularities of σ_0 and σ_1 are Fourier transforms of each other. This is exactly as in §5 of [11] , and (appropriately understood) will be true in the manifold case as well.

Proposition 2.6. *One has the following exact sequence:*

$$0 \to J^{\ell,m-1} \oplus J^{\ell-1,m} \to J^{\ell,m} \overset{\sigma_0}{\to} R^m(\Delta^n \setminus \Sigma) \to 0.$$

Here $R^m(\Delta^n \setminus \Sigma)$ is, roughly speaking, the space of smooth functions on $\Delta^n \setminus \Sigma$ that have singularities of degree m at Σ in the normal directions (for details we refer to [11], §5 and §6).

The classes of operators with kernels in the $J^{\ell,m}$ are closed under composition in the following sense:

Proposition 2.7. *The composition of properly supported operators with kernels in $J^{\ell,m}$ and $J^{\tilde{\ell},\tilde{m}}$, respectively, is an operator with kernel in $J^{\ell+\tilde{\ell}+1/2,m+\tilde{m}-1/2}$. In particular $J^{-1/2,1/2}$ is an algebra:*

$$J^{-1/2,1/2} \circ J^{-1/2,1/2} \subset J^{-1/2,1/2},$$

and for any $u \in J^{\ell,m}$, $v \in J^{\tilde{\ell},\tilde{m}}$, we obtain (ignoring Maslov factors and half-densities):

$$\sigma_0(u \circ v)(\bar{x},\bar{x}) = \sigma_0(u)(\bar{x},\bar{x})\sigma_0(\bar{x},\bar{x}),\ \sigma_1(\bar{x},\bar{y}) = \frac{1}{\sqrt{2\pi}}\int \sigma_1(u)(\bar{x},\bar{z}(t))\sigma_1(\bar{z}(t),\bar{y})dt,$$

where $\bar{z}(t)$ is the bicharacteristic curve joining $\bar{x}$ and $\bar{y}$.

Note that, by the symbol calculus for semiclassical FIOs ([10]) there are formulae for the symbols σ_0, σ_1, of the composition. (Microlocally near $\Delta \setminus \Sigma$ the operators are pseudodifferential and the symbol calculus for σ_0 is the usual one). We will have more to say about the symbol calculus in the next section.

Proof. (Our proof follows the lines of the proof of Theorem 0.1 in [1].) It is not hard to show that the classes J are closed under composition by operators F_1, F_2 as in definition 2.2. Therefore we start with kernels as in (2.1). It is also not hard to show that it suffices to prove the theorem in the case when the amplitudes of these kernels are independent of $\hbar$, in which case $\ell = \tilde{\ell} = -1/2$. Let us therefore consider

$$u(x,y) = \frac{1}{(2\pi\hbar)^n}\int e^{i\hbar^{-1}((x-y)p-sp_1)+is\sigma}a(s,x,y,p,\sigma)dsdpd\sigma,$$

$$v(y,z) = \frac{1}{(2\pi\hbar)^n}\int e^{i\hbar^{-1}((y-z)\tilde{p}-\tilde{s}\tilde{p}_1)+i\tilde{s}\tilde{\sigma}}\tilde{a}(\tilde{s},y,z,\tilde{p},\tilde{\sigma})d\tilde{s}d\tilde{p}d\tilde{\sigma},$$

where a and $\tilde{a}$ are classical symbols in σ and $\tilde{\sigma}$ of orders $m' = m - 1/2$ and $\tilde{m}' = \tilde{m} - 1/2$, respectively. For χ_0 a function that vanishes near the origin, and 1 outside a compact support, we have

$$u(x,y) = \frac{1}{(2\pi\hbar)^n}\int e^{i\hbar^{-1}((x-y)p-sp_1)+is\sigma}a(s,x,y,p,\sigma)\chi_0(\sigma)dsdpd\sigma + \tilde{u},$$

$$v(y,z) = \frac{1}{(2\pi\hbar)^n}\int e^{i\hbar^{-1}((y-z)\tilde{p}-\tilde{s}\tilde{p}_1)+i\tilde{s}\tilde{\sigma}}\tilde{a}(\tilde{s},y,z,\tilde{p},\tilde{\sigma})\chi_0(\tilde{\sigma})d\tilde{s}d\tilde{p}d\tilde{\sigma} + \tilde{v},$$

where $\tilde{u}$, $\tilde{v} \in \text{sc-}I^{\ell}\left(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{F}\partial X^n_c\right)$. We can therefore assume that a, $\tilde{a}$ are zero near $\sigma = 0$, $\tilde{\sigma} = 0$ respectively. The composition has Schwartz kernel:

$$\omega(x,z) = \frac{1}{(2\pi\hbar)^n} \int e^{is\sigma + i\tilde{s}\tilde{\sigma}} D ds d\tilde{s} d\sigma d\tilde{\sigma} dp,$$

with

$$D = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}\phi} \left(a \cdot \tilde{a}\right) \left(s, \tilde{s}, x, y, z, p, \tilde{p}, \sigma, \tilde{\sigma}\right) dy d\tilde{p},$$

where

$$\phi = (x - y)p + (y - z)\tilde{p} - \tilde{s}\tilde{p}_1 - sp_1.$$

The critical points for this phase for each x, z, p fixed are: $\tilde{p} = p$, $y' = z'$, $y_1 = z_1 - \tilde{s}$. So, by the stationary phase theorem, we obtain:

$$D \sim e^{i\hbar^{-1}\phi(y=(z_1-\tilde{s},z'),\tilde{p}=p)} a\Big(s, x, y = (z_1 - \tilde{s}, z'), p, \sigma\Big)\ \tilde{a}\Big(\tilde{s}, y = (z_1 - \tilde{s}, z'), z, p, \tilde{\sigma}\Big)$$

as $\hbar \to 0$. Stationary phase in fact gives us a complete asymptotic expansion in increasing powers of $\hbar$, with coefficients derivatives of a and $\tilde{a}$ evaluated at the critical points. It suffices to consider the contribution to ω of the leading term in the expansion of D (the other terms are treated in the same manner):

$$\omega_0(x,z) := \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(x'-z')p' + (x_1 - z_1 - s - \tilde{s})p_1] + is\sigma + i\tilde{s}\tilde{\sigma}}$$
$$a(s, x, (z_1 - \tilde{s}, z'), p, \sigma)\, \tilde{a}(\tilde{s}, (z_1 - \tilde{s}, z'), z, p, \sigma)\, ds d\tilde{s} dp d\sigma d\tilde{\sigma}.$$

Making the change of variables $t = s + \tilde{s}$, we can write

$$\omega_0(x,z) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(x_1 - z_1 - t)p_1 + (x'-z')p'] + it\tilde{\sigma}} b(t, x, z, p, \tilde{\sigma}) dt dp d\tilde{\sigma},$$

where

$$b(t, x, z, p, \tilde{\sigma}) = \int e^{is(\sigma - \tilde{\sigma})} a(s, x, (z_1 - t + s, z'), p, \sigma)\, \tilde{a}(t - s, (z_1 - t + s, z'), z, p, \tilde{\sigma})\, ds d\sigma.$$

Next, we split ω_0 in three parts, as follows.

Let $\chi_k = \chi_k(\sigma, \tilde{\sigma})$, $k = 1, 2$ be smooth, classical symbols of degree zero such that

$$\chi_1 = \begin{cases} 1 \text{ for } |\sigma| \leq \frac{1}{2}\epsilon|\tilde{\sigma}| \\ \\ 0 \text{ for } |\sigma| \geq \epsilon|\tilde{\sigma}| \end{cases} \quad \text{and} \quad \chi_2 = \begin{cases} 1 \text{ for } |\tilde{\sigma}| \leq \frac{1}{2}\epsilon|\sigma| \\ \\ 0 \text{ for } |\tilde{\sigma}| \geq \epsilon|\sigma| \end{cases}$$

for $\epsilon << 1$. We let

$$\Upsilon_k(x,z) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(x_1 - z_1 - t)p_1 + (x'-z')p'] + it\tilde{\sigma}} b(t, x, z, p, \tilde{\sigma})\, \chi_k\, dt dp d\tilde{\sigma} \quad k = 1, 2,$$

and

$$\Upsilon_0 = \omega_0 - \Upsilon_1 - \Upsilon_2.$$

We will show that Υ_k, $k=1,2$ is a semiclassical state in the flow out while Υ_0 is an integral of the form (2.1).

Let us rewrite, for $k=1,2$,

$$\Upsilon_k = \frac{1}{(2\pi\hbar)^n}\int e^{\hbar^{-1}(x'-z')p'}\left(\int e^{i\hbar^{-1}(x_1-z_1-t)p_1} C_k(x,z,t,p)\,dp_1\,dt\right)\,dp'$$

where

$$C_k(x,z,t,p) := \int e^{it\tilde{\sigma}+is(\sigma-\tilde{\sigma})}\chi_k a\,\tilde{a}\,ds d\sigma d\tilde{\sigma},$$

interpreted as an iterated integral (first with respect to s). On the support of χ_1 one has $|\sigma|\le\epsilon|\tilde{\sigma}|$ which implies $|\sigma-\tilde{\sigma}|\ge|\tilde{\sigma}|-|\sigma|\ge(1-\epsilon)|\tilde{\sigma}|$, and therefore

$$\frac{1}{|\sigma-\tilde{\sigma}|^N}\le\frac{1}{(1-\epsilon)^N}\frac{1}{|\tilde{\sigma}|^N}\le\left(\frac{\epsilon}{1-\epsilon}\right)^N\frac{1}{|\sigma|^N}$$

for all $N>0$. Since $a\ \tilde{a}$ vanish near $\tilde{\sigma}=0=\sigma$ and χ_1 vanishes in a conic neighborhood of the diagonal, we can integrate by parts repeatedly to obtain

$$C_1(x,z,t,p)=\int e^{it\tilde{\sigma}+is(\sigma-\tilde{\sigma})}\frac{\chi_1}{(-i)^N(\sigma-\tilde{\sigma})^N}D_s^N(a\tilde{a})ds d\sigma d\tilde{\sigma}.$$

Since $D_s^N(a\tilde{a})$ is of the same order in $\sigma,\tilde{\sigma}$ and N is arbitrary, C_1 is Schwartz in the variable t. Applying once again the method of stationary phase this implies that

$$\int e^{i\hbar^{-1}(x_1-z_1-t)p_1}C_1(x,z,t,p)\,dp_1\,dt$$

is a semiclassical symbol and therefore Υ_1 is a semiclassical state on the flow out. The proof for Υ_2 is analogous.

To show that Υ_0 is as desired we only need to show that

$$\begin{aligned}b_0(t,x,z,p,\tilde{\sigma}) = \int & e^{is(\sigma-\tilde{\sigma})}a(s,x,(z_1-t+s,z'),p,\sigma)\\ & \tilde{a}(t-s,(z_1-t+s,z'),z,p,\tilde{\sigma})\ (1-\chi_1(\sigma,\tilde{\sigma})-\chi_2(\sigma,\tilde{\sigma}))\ ds d\sigma.\end{aligned}$$

is a classical symbol in $\tilde{\sigma}$ of order $m'+\tilde{m}'$.

Making the change of variables $\tau=\frac{\sigma}{\tilde{\sigma}}$, we obtain:

$$\begin{aligned}b_0 = \tilde{\sigma}\int & e^{is\tilde{(}\tau-1)}a(s,x,(z_1-t+s,z'),p,\tilde{\sigma}\tau)\\ & \tilde{a}(t-s,(z_1-t+s,z'),z,p,\tilde{\sigma})(1-\chi_1(\tilde{\sigma}\tau,\tilde{\sigma})-\chi_2(\tilde{\sigma}\tau,\tilde{\sigma}))ds d\tau.\end{aligned}$$

Notice that τ is bounded in the support of the amplitude, and so using stationary phase in $\tilde{\sigma}$, we get an expansion for b_0 in terms of $\tilde{\sigma}$, where the leading term is

$$b\sim 2\pi a_{m'}(0,x,(z_1-t,z'),p,\tilde{\sigma})\ \tilde{a}_{\tilde{m}'}(t,(z_1-t,z'),z,p,\tilde{\sigma}),$$

and we conclude b is a classical symbol in $\tilde{\sigma}$ of degree $m'+\tilde{m}'$.

The symbol of $\omega(x,z)$ in the diagonal is easily computed as:

$$\sigma_0(\omega)(x,z=x) = 2\pi b_{m'+\tilde{m}'}(t=0,x,z=x,p,p_1=\tilde{\sigma})$$

$$= (2\pi)^2 a_{m'}(0,x,z=x,p,\tilde{\sigma}=p_1)\tilde{a}_{\tilde{m}'}(t=0,z=x,z=x,p,\tilde{\sigma}=p_1)$$

$$= \sigma_0(u)(x,z=x)\sigma_0(v)(z=x,x).$$

The symbol in the flow-out is the sum of the principal symbols of each summand at $(t=z_1-x_1, x'=z', z_1-t+s=x_1+s)$, which after the cancellation of χ_1,χ_2, reduces to

$$\sqrt{2\pi}\int e^{it\tilde{\sigma}}e^{is(\sigma-\tilde{\sigma})}a(s,x,(x_1+s,x'),(0,p'),\sigma)$$
$$\tilde{a}(t-s,(x-1+s,x'),(z_1,x'),(0,p'),\tilde{\sigma})dsd\sigma d\tilde{\sigma}$$

$$= \frac{1}{\sqrt{2\pi}}\int\left(\sqrt{2\pi}\int e^{is\sigma}a(s,x,(x_1+s,x'),(0,p'),\sigma)d\sigma\right)$$
$$\left(\sqrt{2\pi}\int e^{i(t-s)\tilde{\sigma}}\tilde{a}(t-s,(x_1+s,x'),(z_1,x'),(0,p'),\tilde{\sigma})d\tilde{\sigma}\right)ds$$

$$= \frac{1}{\sqrt{2\pi}}\int \sigma_1(u)(x,(x_1+s,x'),(0,p'))\sigma_1(v)((x_1+s,x'),(z_1,x'),(0,p'))ds.$$

□

3. The manifold case

3.1. **The spaces $J^{\ell,m}$ on manifolds.** In this section we extend the definition of the spaces $J^{\ell,m}$ to the manifold case. Let M be a C^∞ manifold of dimension n, and $X_c \subset T^*M$ be a compact domain with smooth boundary contained in the cotangent bundle. The boundary ∂X_c is then foliated by curves tangent to the kernel of the pull-back of the symplectic form. In addition, we assume that the fibrating and Bohr-Sommerfeld conditions are satisfied, i.e., the leaves of the null foliation are the fibers of a submersion $\pi : \partial X_c \to S$, and for each leaf $\gamma \subset \partial X_c$,

$$\int_\gamma \alpha \in 2\pi\mathbb{Z}, \tag{3.1}$$

where α is the tautological one-form in T^*M.

Definition 3.1. Given X_c as above, define the flow-out

$$\mathcal{F}\partial X_c := \left\{(\overline{x},\overline{y}) \;\middle|\; \overline{x},\overline{y}\in\partial X_c,\ \overline{x},\overline{y} \text{ are in the same leaf}\right\} \subset T^*(M\times M). \tag{3.2}$$

The condition that ∂X_c be fibrating easily implies that $\mathcal{F}\partial X$ is a closed submanifold of $T^*(M\times M)$.

Lemma 3.2. *There exists an open neighborhood* $\partial X_c \subset U \subset T^*M$ *of* ∂X_c*, and a map*

$$P : U \to \mathbb{R} \tag{3.3}$$

such that zero is a regular value of P*,* $\partial X_c = P^{-1}(0)$*, the orbits of the Hamilton flow generated by* P *are* 2π *periodic on the boundary* ∂X_c*, and coincide with the leaves of the foliation.*

Remark 3.3. From now on we will fix a defining function of X_c, P, with the properties of this Lemma.

Proof. There exists U and $F : U \to \mathbb{R}$ such that $\partial X_c = F^{-1}(0)$, where zero is a regular value. The Hamiltonian vector field Ξ_F is tangent to $F^{-1}(0) = \partial X_c$, and therefore the trajectories of the Hamiltonian F coincide set-theoretically with the leaves of the foliation. In particular, they are periodic. For each $\overline{x} \in \partial X_c$, let $T(\overline{x})$ denote the minimal period of the trajectory through $\overline{x}$. The fibrating condition can be seen to imply that the function T is smooth. Extend this function to a smooth function $T : U \to \mathbb{R}^+$. The defining function that satisfies the conclusions of the lemma is then

$$P = \frac{T}{2\pi}\, F$$

□

Remark 3.4. Notice that the boundary ∂X_c of X_c may not be connected. In those cases, the flow-out $\mathcal{F}\partial X_c$ consist of the union of Lagrangian that are pairwise disjoint.

Let $\Delta \subset T^*(M \times M)$ be the diagonal in $T^*(M \times M)$.

Lemma 3.5. *The diagonal and the flow-out* $(\Delta, \mathcal{F}\partial X_c)$ *intersect cleanly.*

Proof. Let $(\overline{x}, \overline{x}) \in \Delta \cap \mathcal{F}\partial X_c$. It is easy to show that

$$T_{(\overline{x},\overline{x})}\mathcal{F}\partial X_c = \left\{ (\delta\overline{x} + r\Xi_P(\overline{x}), \delta\overline{x}) \middle| \delta\overline{x} \in T_{\overline{x}}\partial X_c, r \in \mathbb{R} \right\},$$

and so

$$T_{(\overline{x},\overline{x})}\Delta \cap T_{(\overline{x},\overline{x})}\mathcal{F}\partial X_c = \left\{ (\delta\overline{x}, \delta\overline{x}) \middle| \delta\overline{x} \in T_{\overline{x}}\partial X_c \right\} = T_{(\overline{x},\overline{x})}\left(\Delta \cap \mathcal{F}\partial X_c\right)$$

□

By an analogue to Proposition 2.1 in [11], there exists a locally finite covering $\{U_i\}_i$ of $\Delta \cap \mathcal{F}\partial X_c$, $U_i \subset T^*(M \times M)$ open and contractible, where each U_i intersects only one connected component of $\mathcal{F}\partial X_c$, and for each U_i a canonical transformation

$$\chi_i : U_i \longrightarrow T^*\mathbb{R}^{2n}$$

mapping $U_i \cap \Delta$ to Δ^n, and $U_i \cap \mathcal{F}\partial X_c$ to $\mathcal{F}\partial X_c^n$, where $\Delta^n, \mathcal{F}\partial X_c^n$ are the diagonal and flow-out in the model case, respectively.

Definition 3.6. Let $\Sigma = \Delta \cap \mathcal{F}\partial X_c$ be the intersection of the diagonal and the flow-out. The space $J^{\ell,m}(M \times M; \Delta, \mathcal{F}\partial X_c)$ is the set of families of functions $A(x, y, \hbar)$ which can be written in the form

$$A = A_0 + A_1 + \sum \omega_i$$

where:

(1) $A_0 \in \text{sc}{-}I^{\ell+m}(M \times M; \Delta \setminus \Sigma)$ and $A_1 \in \text{sc}{-}I^{\ell}(M \times M; \mathcal{F}\partial X_c)$, and
(2) ω_i is microlocally supported in U_i and is of the form

$$\omega_i = F_i(v_i),$$

where $v_i \in J^{\ell,m}(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n, \mathcal{F}\partial X_c^n)$ and F_i is a semiclassical Fourier integral operator associated to χ_i^{-1}.

Remark 3.7. As in [11], one can show that the definition does not depend on the choice of the semiclassical FIOs F_i.

Proposition 3.8. *In the general case the conclusion of Proposition 2.4 still holds, namely: Operators with kernel* $A \in J^{\ell,m}(M \times M;\ \Delta, \mathcal{F}\partial X_c)$ *have frequency set contained in the union*

$$FS(A) \subset \Delta' \cup \mathcal{F}\partial X_c'.$$

Away from the intersection Σ *they are microlocally semiclassical Fourier integral operators* $sc{-}I^{\ell+m}(M \times M; \Delta \setminus \Sigma)$ *and* $sc{-}I^{\ell}(M \times M, \mathcal{F}\partial X_c \setminus \Sigma)$ *respectively. Furthermore, one has well-defined symbol maps (ignoring Maslov factors)*

$$\begin{tikzcd} & {|\wedge|^{1/2}(\Delta \setminus \Sigma)} \\ {J^{\ell,m}(M \times M;\ \Delta, \mathcal{F}\partial X_c)} \arrow[ur, "\sigma_0"] \arrow[dr, "\sigma_1"'] & \\ & {|\wedge|^{1/2}(\mathcal{F}\partial X_c \setminus \Sigma).} \end{tikzcd} \tag{3.4}$$

The proof of the existence of the symbol calculus is the subject of the following section.

Our construction, in particular, gives a way to associating to X_c an algebra of operators, which can be thought of as a quantization of X_c:

Definition 3.9. We will denote by $\mathcal{A}_{X_c}$ the space of elements in $J(M \times M; \Delta, \partial X_c)$ that are microlocally of order $\mathrm{O}(\hbar^{\infty})$ in the complement of X_c.

It is not hard to see that $\mathcal{A}_{X_c}$ is indeed closed under composition. (Elements in this algebra correspond to amplitudes that are of order $-\infty$ in σ as $\sigma \to -\infty$.)

3.2. **Symbolic calculus.** Here we discuss how Proposition 3.8 can be proved (for $\hbar$ tending to zero along certain sequences) using the methods from [23]. We begin by recalling the ideas and results from *op. cit.* that we will need.

The pre-quantum circle bundle of T^*M can be identified with the following submanifold of $T^*(M \times S^1)$:

$$Z = \{(x, \theta; \xi, \kappa) \in T^*(M \times S^1)\ ;\ \kappa = 1\},$$

with the obvious circle action. The connection form, α, is the pull-back to Z of the canonical one form of $T^*(M \times S^1)$.

Definition 3.10. ([23]) A Lagrangian submanifold $\Lambda \subset T^*M$ will be called *admissible* iff there exists a conic Lagrangian submanifold,

$$\tilde{\Lambda} \subset T^*(M \times S^1) \cap \{\kappa > 0\}$$

such that

$$\Lambda = \left(\tilde{\Lambda} \cap Z\right)/S^1. \tag{3.5}$$

We call such a $\tilde{\Lambda}$ a *homogenization* of Λ.

It is not hard to see that Λ is admissible if and only if it satisfies the following Bohr-Sommerfeld condition: There exists $\phi : \Lambda \to S^1$ such that

$$\iota^*\eta = d \log \phi,$$

where $\iota : \Lambda \to T^*M$ is the inclusion and η the canonical one form of T^*M. Given such a ϕ, a homogenization of Λ can be defined by:

$$\tilde{\Lambda} = \{e^{i\theta} = \phi(\lambda),\ \kappa > 0\}. \tag{3.6}$$

Definition 3.11. Let M be a C^∞ manifold and consider a family of smooth functions $\{\psi_\hbar\}$. The $\hbar$ *transform* of the family $\psi_\hbar$ is the following distribution (if the series converges weakly) in $M \times S^1$:

$$\Psi(x, \theta) = \sum_{m=0}^{\infty} \psi_{1/m}(x) e^{im\theta}.$$

The main point of the previous two definitions is the following

Lemma 3.12. *([23]) The $\hbar$ transform of a semiclassical state associated to an admissible Lagrangian is a Lagrangian distribution associated to a homogenization of the Lagrangian submanifold.*

We claim that the previous lemma generalizes to our spaces $J^{\ell,m}$ of semiclassical pseudodifferential operators with singular symbols, so that the $\hbar$ transform of their kernels are Guillemin-Uhlmann operators (i. e. those defined in [11]) on $M \times S^1$.

Thus we now consider Δ to be the diagonal in $T^*(M\times M)$ and $\mathcal{F}\partial X_c$ as in (3.2). Clearly a homogenization for Δ is the diagonal $\widetilde{\Delta}$ of $T^*\left(M\times S^1\right)^+ = T^*\left(M\times S^1\right)\cap\{\kappa>0\}$. Let P be a globally defining function for ∂X_c with periodic Hamilton flow on it, as in (3.3). Then a homogenization for $\mathcal{F}\partial X_c$ is the flow-out of the homogenization of P, which is the function $\tilde{P} : T^*\left(U\times S^1\right)^+ \to \mathbb{R}$ defined as

$$\widetilde{P}(x,e^{i\theta};\xi,\kappa)=\kappa P(x,\xi/\kappa),$$

where U is the neighborhood of ∂X_c described in Lemma 3.2. Specifically,

$$\widetilde{\mathcal{F}\partial X_c} = \left\{ \left(\Phi_s^{\widetilde{P}}(x,e^{i\theta},\xi,\kappa)\ ;\ (x,e^{i\theta},\xi,\kappa)\right) \mid s\in\mathbb{R},\kappa>0,\ P(x,\xi/\kappa)=0\right\} \tag{3.7}$$

where $\Phi_s^{\widetilde{P}}$ is the Hamilton flow generated by the equations (here $p=\xi/\kappa$)

$$\dot{x}=\tfrac{\partial P}{\partial p}(x,p), \qquad \dot{\theta}=-p\ \tfrac{\partial P}{\partial p}(x,p)$$

$$\dot{\xi}=-\kappa\tfrac{\partial P}{\partial x}(x,p), \qquad \dot{\kappa}=0.$$

Notice that $\widetilde{\mathcal{F}\partial X_c}$ is closed if the flow Φ_t^P is 2π periodic, which happens if the Bohr-Sommerfeld condition

$$\int_\gamma pdx\in 2\pi\mathbb{Z}$$

is satisfied for orbits $\gamma\subset\partial X_c$ of P. The function $\phi:\mathcal{F}\partial X_c\to S^1$ associated to this homogenization is

$$\phi(\overline{x},\overline{y})=e^{-i\int_{\overline{x}}^{\overline{y}}pdx},$$

where $\int_{\overline{x}}^{\overline{y}}pdx$ is the action from $\overline{x}$ to $\overline{y}$, and the integral is taken on the curve in the leaf joining $\overline{x}$ to $\overline{y}$.

3.3. **The existence of the symbolic calculus.** Notice that Δ and $\mathcal{F}\partial X_c$ intersect cleanly. As a result, their homogenizations $\widetilde{\Delta}$ and $\widetilde{\mathcal{F}\partial X_c}$ are conic and intersect cleanly too, forming an intersecting pair, in the sense of [22, 11]. The relationship between the semiclassical objects defined above and the operators described in [11] is as follows:

Theorem 3.13. *Let $\Delta,\mathcal{F}\partial X_c$ be as above, and $\widetilde{\Delta},\widetilde{\mathcal{F}\partial X_c}$ their homogenization, respectively. Then any operator $A\in J^{\ell,m}(M\times M\ ;\ \Delta,\mathcal{F}\partial X_c)$ if and only if its $\hbar$-Transform belongs to $I^{\ell,m}\left(M\times S^1\times M\times S^1\ ;\ \widetilde{\Delta},\widetilde{\mathcal{F}\partial X_c}\right)$.*

Proof. We sketch the ideas of the proof in the model case. Let $A(x,y,\hbar)$ be as in equation (2.1). Let us assume first that the amplitude $\frac{1}{(2\pi\hbar)^n}a$ does not depend on $\hbar$ so that

$$A(x,y,\hbar)=\int e^{i\hbar^{-1}[(x_1-y_1-s)p_1-(x'-y')p']+is\sigma}a(s,x,y,p,\sigma)dsdpd\sigma,$$

where $a(s,x,y,p,\sigma)$ is a classical symbol in σ of order $m' = m-1/2$, and compactly supported in s,x,y,p. The $\hbar$-transform of A is then

$$A(x,\theta,y,\alpha) = \sum_{k=1}^{\infty} \int e^{ik[(x_1-y_1-s)p_1+(x'-y')p'+(\theta-\alpha)]+is\sigma} a(s,x,y,p,\sigma)dsdpd\sigma$$

$$= \int \frac{e^{i[(x_1-y_1-s)p_1+(x'-y')p'+(\theta-\alpha)]}}{1-e^{i[(x_1-y_1-s)p_1+(x'-y')p'+(\theta-\alpha)]}} e^{is\sigma} a(s,x,y,p,\sigma)dsdpd\sigma.$$

This distribution is the push-forward under the projection $(s,x,y,\theta,\alpha,p) \to (x,y,\theta,\alpha)$ of the product of the distributions

$$\Gamma(s,x,y,\theta,\alpha,p) = \frac{e^{i[(x_1-y_1-s)p_1+(x'-y')p'+(\theta-\alpha)]}}{1-e^{i[(x_1-y_1-s)p_1+(x'-y')p'+(\theta-\alpha)]}}$$

and

$$\Upsilon(s,x,y,\theta,\alpha,p) = \int e^{is\sigma} a(s,x,y,p,\sigma)d\sigma.$$

Γ is a distribution in space conormal to the hypersurface $(x_1-y_1-s)p_1+(x'-y')p' = -(\theta-\alpha)$, while Υ is a distribution conormal to $s=0$. Therefore, $A(x,\theta,y,\alpha)$ is a Guillemin-Uhlmann distribution associated to the pair $\left(\widetilde{\Delta^n}, \widetilde{\mathcal{F}\partial X_c^n}\right)$. The general case (i.e. when a also depends on $\hbar$) is an asymptotic sum of derivatives and integrals (with respect to the θ variables) of the previous case. The converse is also true, and the proof is analogue to that in [23] for semiclassical states. $\square$

This proposition together with Proposition 2.4 of [23], relating the frequency set of an $\hbar$-dependent vector and the wave-front set of its $\hbar$ transform, implies part (1) of Theorem 1.1.

The symbols of operators in $J^{\ell,m}(\Delta, \mathcal{F}\partial X_c)$ are the reduction of the symbols of its $\hbar$-Transform, in the following sense. Let $\overline{x}, \overline{y} \in \mathcal{F}\partial X_c \setminus \Sigma$, $\overline{x} = \Phi_s^P(\overline{y})$ for some $s \in \mathbb{R}$, and take

$$(\widetilde{x},\widetilde{y}) = \left(\Phi_s^{\widetilde{P}}(\overline{y}, e^{i\theta=0}=1, \kappa=1), (\overline{y}, e^{i\theta=0}=1, \kappa=1)\right) \in \widetilde{\mathcal{F}\partial X_c^n}.$$

Using (3.7), we obtain an isomorphism between $T_{(\widetilde{x},\widetilde{y})}\widetilde{\mathcal{F}\partial X_c}$ and $T_{(\overline{x},\overline{y})}\mathcal{F}\partial X_c \times \mathbb{R} \times \mathbb{R}$, which leads to

$$\left|T_{(\widetilde{x},\widetilde{y})}\widetilde{\mathcal{F}\partial X_c}\right|^{1/2} \cong \left|T_{(\overline{x},\overline{y})}\mathcal{F}\partial X_c \times \mathbb{R} \times \mathbb{R}\right|^{1/2} \cong \left|T_{(\overline{x},\overline{y})}\mathcal{F}\partial X_c\right|^{1/2} \tag{3.8}$$

Therefore, every half-density in $T_{(\widetilde{x},\widetilde{y})}\widetilde{\mathcal{F}\partial X_c}$ will define a half-density in $T_{(\overline{x},\overline{y})}\mathcal{F}\partial X_c$. Let $\widetilde{\Sigma} = \widetilde{\mathcal{F}\partial X_c} \cap \widetilde{\Delta}$. For each family $A \in J^{\ell,m}$, denote its $\hbar$-Transform by $\widetilde{A}$. There is a well defined symbol map

$$\widetilde{\sigma_1} = \sigma_1\left(\widetilde{A}\right)_{\widetilde{\mathcal{F}\partial X_c}-\widetilde{\Sigma}} \in C^\infty\left(\widetilde{\mathcal{F}\partial X_c} \setminus \widetilde{\Sigma}, \widetilde{\Omega_1} \bigotimes \widetilde{L_1}\right),$$

where $\widetilde{\Omega}_j$ is the bundle of half-densities on $\widetilde{\mathcal{F}\partial X_c}$, and $\widetilde{L}_i$ is the corresponding Maslov bundle. Ignoring Maslov factors, we can define the symbols on $\mathcal{F}\partial X_c \setminus \Sigma$ by the identification (3.8), i.e., restricting the symbol of the $\hbar$-Transform to $\theta = 0$, $\kappa = 1$ ($\sigma_1 = \widetilde{\sigma_1}\big|_{\theta=0,\kappa=1}$). The construction of the principal symbol on the diagonal is similar.

Let $\Sigma^n = \Delta^n \cap \mathcal{F}\partial X_c^n$, $\widetilde{\Sigma}^n = \widetilde{\Delta^n} \cap \widetilde{\mathcal{F}\partial X_c^n}$. In the model case, we know by [11] that $A(x, \theta, y, \alpha)$ is microlocally in

$$I^{\ell+m}\left(\mathbb{R}^n \times S^1 \times \mathbb{R}^n \times S^1\ ;\ \widetilde{\Delta^n} \setminus \widetilde{\Sigma}^n\right),\ \text{and}\ I^{\ell}\left(\mathbb{R}^n \times S^1 \times \mathbb{R}^n \times S^1\ ;\ \widetilde{\mathcal{F}\partial X_c^n} \setminus \widetilde{\Sigma}^n\right).$$

Therefore $A_\hbar$ is microlocally in sc-$I^{\ell+m}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n \setminus \Sigma^n\right)$ and sc-$I^{\ell}\left(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{F}\partial X_c^n \setminus \Sigma^n\right)$ in the sense that the $\hbar$-transform is microlocally in their corresponding spaces. We have proved the following

Proposition 3.14.

$$\cap_\ell J^{\ell,m} = sc\text{-}I^{\infty}\left(\mathbb{R}^n \times \mathbb{R}^n; \Delta^n\right)$$

and

$$\cap_m J^{\ell,m} = sc\text{-}I^{\ell}\left(\mathbb{R}^n \times \mathbb{R}^n; \mathcal{F}\partial X_c^n\right).$$

3.4. **A symbolic compatibility condition.** Recall that the foliation of ∂X_c is fibrating, i. e., there exists a C^∞ Hausdorff manifold S and a smooth fiber map

$$\pi : \partial X_c \to S, \tag{3.9}$$

whose fibers are the connected leaves of the foliation defined Section 3.1. Elements of $\mathcal{F}\partial X_c$ are pairs of points in ∂X_c that lie in the same fiber of π.

Generalizing a construction in [15], given $s \in S$ let $\mathfrak{SO}_s$ be the $*-$algebra of all pseudodifferential operators, acting on the space of half-densities $C^\infty\left(|F_s|^{1/2}\right)$, where F_s is the fiber of $\pi : \partial X_c \to S$ above s. Let $\mathfrak{SO}$ be the sheaf of $*-$algebras on S whose stalk at s is $\mathfrak{SO}_s$. We will say that a section k of $\mathfrak{SO}$ is smooth if the Schwartz kernel of the operator $k(s)$ depends smoothly on $s \in S$. The Schwartz kernel theorem, applied fiber-wise to the fibers of π, together with the natural symplectic structure of S yield the following:

Proposition 3.15. *The vector space of smooth sections of the sheaf $\mathfrak{SO}$ is naturally isomorphic to the space of half-densities on $\mathcal{F}\partial X_c \setminus \Sigma$ that extend to $\mathcal{F}\partial X_c$ as a conormal distribution to Σ*

Proof. Let $\gamma = (\gamma_1, \gamma_2) \in \mathcal{F}\partial X_c$, $s = \pi(\gamma_1) = \pi(\gamma_2) \in S$. One then has the following fiber product diagram:

$$\begin{tikzcd}
T_\gamma \mathcal{F}\partial X_c \arrow[r, "d\pi_1"] \arrow[d, "d\pi_2"'] & T_{\gamma_1}\partial X_c \arrow[d, "d\pi"] \\
T_{\gamma_2}\partial X_c \arrow[r, "d\pi"] & T_s S
\end{tikzcd}$$

and so we get the exact sequence

$$\begin{array}{c}
0 \longrightarrow T_\gamma \mathcal{F}\partial X_c \longrightarrow T_{\gamma_1}\partial X_c \oplus T_{\gamma_2}\partial X_c \longrightarrow T_s S \longrightarrow 0 \\
v \longmapsto (d\pi_1 v, d\pi_2 v) \\
(v_1, v_2) \longmapsto d\pi(v_2) - d\pi(v_1)
\end{array}$$

This gives a natural identification

$$\mathbb{C} \cong |T_\gamma \mathcal{F}\partial X_c|^{1/2} \otimes |T_{\gamma_1}\partial X_c \oplus T_{\gamma_2}\partial X_c|^{-1/2} \otimes |T_s S|^{1/2}.$$

Since S is a symplectic manifold, there is a canonical half-density on $T_s S$, and we get an identification

$$\left|T_\gamma \mathcal{F}\partial X_c\right|^{1/2} \cong \left|T_{\gamma_1}\partial X_c \oplus T_{\gamma_2}\partial X_c\right|^{1/2} \cong \left|T_{\gamma_1}\partial X_c\right|^{1/2} \otimes \left|T_{\gamma_2}\partial X_c\right|^{1/2}.$$

Finally, given a half density in $T_{\gamma_k}\partial X_c$, $k = 1, 2$, we need to get a half-density in $T_{\gamma_k} F_s$, where F_s is the fiber of π above s. We have the following exact sequence:

$$0 \longrightarrow \ker\ d\pi_{\gamma_k} \longrightarrow T_{\gamma_k}\partial X_c \longrightarrow T_s S \longrightarrow 0$$

which, by the same process as above, gives an identification

$$\left|T_{\gamma_k} F_s\right|^{1/2} = \left|\ker\ d\pi_{\gamma_k}\right|^{1/2} \cong \left|T_{\gamma_k}\partial X_c\right|^{1/2}$$

Hence

$$\left|T_\gamma \mathcal{F}\partial X_c\right|^{1/2} \cong \left|T_{\gamma_1} F_s\right|^{1/2} \otimes \left|T_{\gamma_2} F_s\right|^{1/2} \cong \left|T_{(\gamma_1,\gamma_2)}(F_s \times F_s)\right|^{1/2}$$

This gives a smooth isomorphism between two line bundles over $\mathcal{F}\partial X_c$: $\left|T\mathcal{F}\partial X_c\right|^{1/2}$ and the line bundle $\Upsilon \to \mathcal{F}\partial X_c$ whose fiber over (γ_1, γ_2) is $\left|T_{(\gamma_1,\gamma_2)}(F_s \times F_s)\right|^{1/2}$, where $\pi(\gamma_1) = s = \pi(\gamma_2)$. Clearly a section of the sheaf $\mathfrak{G}\mathfrak{O}$ is a distributional section of Υ conormal to Σ, and by the previous isomorphism this is equivalent to a distributional section of $\left|T\mathcal{F}\partial X_c\right|^{1/2}$ conormal to Σ. □

The previous isomorphism yields an algebra structure on the space of distributional half densities on $\mathcal{F}\partial X_c$ which are conormal to Σ. Analogously as in [15] (Proposition 2.7), one can see that the algebraic structure on this space is given, away from Σ, by the composition of half densities regarded as symbols of Fourier integral operators associated to $\mathcal{F}\partial X_c$.

Let us now take $A \in J^{\ell,m}$. The symbol $\sigma_1(A)$ in $\mathcal{F}\partial X_c \setminus \Sigma$ blows-up as the point where σ_1 is evaluated tends to the intersection. In fact, in [1] it was shown that this symbol has a natural extension to a distribution conormal to Σ. Using the same identification above, this determines the kernel of a Pseudodifferential operator on the fiber above each point of S. We have proved:

Proposition 3.16. *For $A \in J^{\ell,m}(M \times M, \Delta, \mathcal{F}\partial X_c)$, the symbol $\sigma_1(A)$ can be identified with a global section of the sheaf $\mathfrak{S}\mathfrak{O}$, that is, with a family of classical pseudodifferential operators of order $m' = m - 1/2$ acting on fibers of $\pi : \partial X_c \to S$ (orbits in the flow-out).*

For each $s \in S$ and F_s the corresponding fiber above s, let us denote this operator by $\sigma_1(A)_s$:

$$C^\infty(F_s) \xrightarrow{\sigma_1(A)_s} C^\infty(F_s)$$

As a result, there is a well-defined symbol

$$\sigma(\sigma(A)_s) : T^*F_s \setminus 0 \to \mathbb{C}.$$

As we will now see, this symbol is related to the symbol $\sigma_0(A)$ of A on the diagonal. (This is the compatibility of the symbols of A announced earlier.)

Let P be a defining function of ∂X_c with a 2π-periodic flow on ∂X_c. We have the following diffeomorphism

$$\begin{tikzcd} \partial X_c \times S^1 \arrow[r] & \mathcal{F}\partial X_c \\ (\overline{x}, t) \arrow[r, mapsto] & \left(\Phi_t^P \overline{x}, \overline{x}\right) \end{tikzcd}$$

from which one can see that, for any $\gamma = (\overline{x}, \overline{x}) \in \Sigma = \Delta \cap \mathcal{F}\partial X_c$, there is a natural isomorphism

$$N_\Sigma^{\mathcal{F}\partial X_c} := T_\gamma \mathcal{F}\partial X_c / T_\gamma \Sigma \cong T_{\overline{x}} F_s$$

($N_\Sigma^{\mathcal{F}\partial X_c}$ is the normal space to Σ in $\mathcal{F}\partial X_c$ at γ). Therefore, for each $\overline{x} \in \partial X_c$, $T^*_{\overline{x}} F_s$ is isomorphic to the dual space $\left(N_\Sigma^{\mathcal{F}\partial X_c}\right)^*$.

Now let $N_\Sigma^\Delta := T_\gamma \Delta / T_\gamma \Sigma$ be the normal space to Σ in Δ at γ. By [11], $N_\Sigma^{\mathcal{F}\partial X_c}$ and N_Σ^Δ are supplementary Lagrangian subspaces of the two-dimensional symplectic vector space $W = (T_\gamma \Sigma)^\perp / T_\gamma \Sigma$. Therefore, $N_\Sigma^{\mathcal{F}\partial X_c}$ and N_Σ^Δ are in duality with each other (they are canonically paired by the symplectic form of W). In the end we obtain a natural isomorphism

$$T^*_{\overline{x}} F_s \cong \left(N_\Sigma^{\mathcal{F}\partial X_c}\right)^* \cong N_\Sigma^\Delta.$$

The symbol of A on the diagonal belongs to a class $S^{m'}(\Omega_0; \Delta, \Sigma)$ of smooth functions on $\Delta \setminus \Sigma$ which blow up at a prescribed rate at Σ (see [11] for more details). Every element in this class determines a smooth function on $N^{\Delta}_{\Sigma} \setminus \{0\}$. The *compatibility condition* alluded to in the Introduction is the following:

Theorem 3.17. *The symbol of* $A \in J^{\ell,m}(M \times M, \Delta, \mathcal{F}\partial X_c)$ *on the flow-out, identified with a family* $\{\sigma_1(A)_s\}_{s\in S}$ *of pseudodifferential operators along the fibers* F_s*, satisfies that for each* $\overline{x} \in F_s$

$$\begin{aligned}&\sigma(\sigma_1(A)_s)_{\overline{x}}(\tau) = \lim_{\substack{\overline{y}\to\overline{x}\\ \overline{y}\in X_c\setminus\partial X_c}} \tfrac{\sigma_0(A)(\overline{y},\overline{y})}{P^{m'}(\overline{y})}\\ &\textit{and}\\ &\sigma(\sigma_1(A)_s)_{\overline{x}}(-\tau) = \lim_{\substack{\overline{y}\to\overline{x}\\ \overline{y}\in T^*M\setminus X_c}} \tfrac{\sigma_0(A)(\overline{y},\overline{y})}{P^{m'}(\overline{y})},\end{aligned} \tag{3.10}$$

where $\tau \in T^*_{\overline{x}}F_s$ *is the dual of the Hamilton field of* P *at* $\overline{x}$ *regarded as an element in* $T_{\overline{x}}F_s$*, and the two limits are taken from the interior of* X_c *and the exterior of* X_c*, respectively. Moreover, this condition is intrinsic, i.e., it does not depend on the choice of* P*.*

The proof reduces to the model case, where it is immediate. One can also verify directly that changing P by a multiplicative factor does not alter the relationships (3.10).

The symbols in the flow-out become more natural under the present setting, as can be seen in the following symbolic version of Proposition 2.7.

Proposition 3.18. *The composition of properly supported operators with kernels in* $J^{\ell,m}$ *and* $J^{\tilde{\ell},\tilde{m}}$*, respectively, is an operator with kernel in* $J^{\ell+\tilde{\ell}+1/2,m+\tilde{m}-1/2}$*. For any* $A \in J^{\ell,m}$*,* $B \in J^{\tilde{\ell},\tilde{m}}$*, we obtain the usual symbol in the diagonal:*

$$\forall \overline{x} \in X_c \setminus \partial X_c \quad \sigma_0(A \circ B)(\bar{x},\bar{x}) = \sigma_0(A)(\bar{x},\bar{x})\sigma_0(B)(\bar{x},\bar{x}),$$

and for any fiber F_s *above* $s \in S$*,*

$$\sigma_1(A \circ B)_s = \sigma_1(A)_s \circ \sigma_1(B)_s\ .$$

3.5. **The adjoint.** The class $J^{\ell,m}$ is closed under the operation of taking adjoints, and information about the symbol of the adjoint is given in the following

Proposition 3.19. *Let* $A \in J^{\ell,m}(M \times M; \Delta, \mathcal{F}\partial X_c)$*, then the adjoint* A^* *belongs again to* $J^{\ell,m}(M \times M; \Delta, \mathcal{F}\partial X_c)$*, and*

$$\sigma_0(A^*)(\overline{x},\overline{x}) = \overline{\sigma_0(A)(\overline{x},\overline{x})} \quad \textit{for } (\overline{x},\overline{x}) \in \Delta \setminus \Sigma, \textit{ and}$$

$$\sigma_1(A^*)_s = (\sigma_1(A)_s)^*\,, \quad \textit{for each } s \in S.$$

Proof. The first statement is as in the usual theory of $\hbar$-pseudodifferential operators. It is enough to prove the rest in the model case. Take A with Schwartz kernel

$$K(x,y,h) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}((x-y)\cdot p - sp_1) + is\sigma} a(s,x,y,p,\sigma,\hbar) ds dp d\sigma,$$

where $a \in \mathcal{A}^{\ell',m'}_{classical}$, $\ell' = \ell + 1/2$, $m' = m - 1/2$. The Schwartz kernel of the adjoint is given by

$$K^*(x,y) = \overline{K(y,x)} = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}((x-y)p - sp_1) + is\sigma} \overline{a(-s,y,x,p,\sigma,\hbar)} ds dp d\sigma,$$

where $a(s,x,p,\sigma,\hbar)$ was replaced by $\overline{a(-s,y,x,p,\sigma,\hbar)}$. Taking $\overline{x} = (x,(p_1 = 0,p')), \overline{y} = \phi^P_s \overline{x} = ((x_1+s,x'),(p_1=0,p'))$, the symbol in the flow-out is given by

$$\sigma_1(A^*)(\phi^P_s\overline{x},\overline{x}) = \sqrt{2\pi} \int \overline{a_{-\ell'}(-s,y,x,p,\sigma)} e^{is\sigma} d\sigma \sqrt{dx dy_1 dp'}\Big|_{x'=y',p_1=0,s=x_1-y_1}$$

$$= \sqrt{2\pi}\overline{\int a_{-\ell'}(-s,y,x,p,\sigma) e^{-is\sigma} d\sigma} \sqrt{dx dy_1 dp'}\Big|_{x'=y',p_1=0,y_1-x_1=-2} = \overline{\sigma_1(A)(\overline{x},\phi^P_s\overline{x})}$$

The proof is now clear, since the symbol $\sigma_1(A^*)$ intertwines the variables and takes the conjugate of $\sigma_1(A)$. □

4. Asymptotics of the trace

4.1. **The trace in case $m' \geq 0$.** We now assume that X_c is compact and ∂X_c is fibrating, as in section 3. We will prove:

Theorem 4.1. *Let $\widehat{A}$ be an operator in the class $\mathcal{A}_{X_c}$. Then, if $m > 1/2$,*

$$\mathrm{Tr}(\widehat{A}) = (2\pi)^{-n}\hbar^{-\ell-m-n} \int_{X_c} \sigma_0(x,x,p,-p) dx dp + O(\hbar^{-\ell-m-n+1}), \tag{4.1}$$

where σ_0 is the symbol of A on the diagonal. If $m = 1/2$,

$$\mathrm{Tr}(\widehat{A}) = (2\pi)^{-n}\hbar^{-\ell-m-n} \int_{X_c} \sigma_0(x,x,p,-p) dx dp + O(\hbar^{-\ell-m-n}\ \hbar\ \log(1/\hbar)). \tag{4.2}$$

The rest of this subsection is devoted to a proof of this result, which we break into a series of lemmas. Note that it is enough to estimate the integral along the diagonal of the kernel of A in the model case with $\ell' = 0$. Consider therefore a semiclassical kernel of the form:

$$u_\hbar(x,y) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(x_1-y_1-s)p_1 + (x'-y')p'] + is\sigma} a(s,x,y,p,\sigma) ds dp d\sigma, \tag{4.3}$$

where $p = (p_1,p')$, $x = (x_1,x')$, $y = (y_1,y')$ and a is a classical symbol in σ of degree $m' = m - 1/2$, compactly supported in (s,x,y,p).

Lemma 4.2. *Let* $\mu_0 > 0$ *be large enough so that* $a|_{\{|p_1|>\mu_0/2\}} = 0$*, and let* $\rho \in C_0^\infty$ *be a smooth function with compact support which is equal to one in* $[-\mu_0/2, \mu_0/2]$ *and is supported in* $[-\mu_0, \mu_0]$*. Then*

$$u_\hbar(x,x) = \frac{1}{(2\pi\hbar)^n} \int e^{-isp_1/\hbar + is\sigma} a(s,x,x,p,\sigma)\rho(\hbar\sigma)\,ds\,dp\,d\sigma + O(\hbar^\infty), \tag{4.4}$$

uniformly in x*.*

Proof. The rigorous definition of u_h when the integral in σ diverges is given in equation (2.3): If $K >> 0$ ($K \geq m'/2 + 1$), then u_h is equal to the absolutely convergent integral

$$\frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(x_1-y_1-s)p_1+(x'-y')p']+is\sigma} \frac{1}{(1+\sigma^2)^K} D^K \left[e^{-is\hbar^{-1}p_1} a(s,x,y,p,\sigma)\right] ds\,dp\,d\sigma,$$

and therefore

$$u_\hbar(x,y) = \frac{1}{(2\pi\hbar)^n} \sum_{j=0}^{K} \int e^{i\hbar^{-1}[(x-y)p-sp_1]+is\sigma} \left(-i\hbar^{-1}p_1\right)^{2K-j} \frac{a_j(s,x,y,p,\sigma)}{(1+\sigma^2)^K} ds\,dp\,d\sigma,$$

where the last expression is obtained by expanding the action of D^K. Note that, $\forall j$, a_j consists of linear combinations of derivatives of a with respect to s (and therefore $a_j \in \mathcal{A}^{0,m'}$). Using this we get that the remainder in equation (4.4) is equal to

$$\sum_{j=0}^{K} \frac{1}{(2\pi\hbar)^n} \int e^{-isp_1/\hbar + is\sigma} \frac{a_j(s,x,x,p,\sigma)}{(1+\sigma^2)^K} \left(-i\hbar^{-1}p_1\right)^{2K-j} (1-\rho(\hbar\sigma))\,ds\,dp\,d\sigma.$$

Let $b_j(s,x,p,\sigma\ ;\ \hbar) = \frac{a_j(s,x,x,p,\sigma)}{(1+\sigma^2)^K}(-i\hbar^{-1}p_1)^{2K-j}(1-\rho(\hbar\sigma))$. We will show that for each j,

$$B_j := \frac{1}{(2\pi\hbar)^n} \int e^{-isp_1\hbar^{-1}+is\sigma} b_j\,ds\,dp\,d\sigma$$

is $O(\hbar^\infty)$

Starting with the change of variables $\mu = \hbar\sigma$, $\omega = -p_1 + \mu$, we obtain that

$$\begin{aligned} B_j &= \frac{\hbar^{-1}}{(2\pi\hbar)^n} \int e^{is\omega/\hbar}\, b_j(s,x,(-\omega+\mu,p'),\hbar^{-1}\mu\ ;\hbar)\,ds\,d\omega\,dp'\,d\mu \\ &= \frac{1}{(2\pi\hbar)^n} \int e^{-i\hbar\xi_1\xi_2}\, c_j(x,\mu,p'\ ;\ \hbar\ ;\ \xi)\,d\xi\,dp'\,d\mu, \end{aligned}$$

where $\xi = (\xi_1, \xi_2)$ are the dual variables to (s,ω), and

$$c_j(x,\mu,p'\ ;\ \hbar\ ;\ \xi) = \frac{1}{2\pi} \int e^{-i(s,\omega)\cdot\xi}\, b_j(s,x,(-\omega+\mu,p'),\hbar^{-1}\mu\ ;\ \hbar)\,ds\,d\omega$$

is the Fourier transform of b_j in the (s,ω) variables. Using the inequality

$$\left| e^{it} - \sum_{k=0}^{N-1} \frac{(it)^k}{k!} \right| \le \frac{|t|^N}{N!}, \tag{4.5}$$

we obtain that for each $N > 0$,

$$\int e^{-i\hbar\xi_1\xi_2} c_j(x,\mu,p'\ ;\ \hbar\ ;\ \xi)d\xi - \sum_{k=0}^{N-1} \int \frac{(-i\hbar\xi_1\xi_2)^k}{k!} c_j(x,\mu,p'\ ;\ \hbar\ ;\ \xi)d\xi$$

$$= \int R_N(\xi\ ;\ \hbar) c_j d\xi,$$

where $|R_N(\xi\ ;\ \hbar)| \le \frac{\hbar^N|\xi_1\xi_2|^N}{N!}$. Moreover, for each $k = 0,\dots,N-1$,

$$\frac{(i\hbar)^k}{k!} \int (-\xi_1\xi_2)^k c_j d\xi = \frac{2\pi(-i\hbar)^k}{k!} \left(\frac{\partial^2}{\partial s\partial p_1}\right)^k b_j(s,x,p,\hbar^{-1}\mu\ ;\ \hbar)\big|_{s=0,p_1=\mu}$$

$$= \frac{2\pi(-i\hbar)^k\ (1-\rho(\mu))}{(1+(\hbar^{-1}\mu)^2)k!} \left(\frac{\partial^2}{\partial s\partial p_1}\right)^k a_j(s,x,(\mu,p'),\hbar^{-1}\mu)\big|_{s=0,p_1=\mu} = 0,$$

since $\rho(\mu)$ is equal to one in the support of a. It follows that

$$\left| \int e^{-i\hbar\xi_1\xi_2} c_j d\xi \right| \le \frac{\hbar^N}{N!} \int \left| (\xi_1\xi_2)^N c_j \right| d\xi$$

$$= \frac{\hbar^N}{2\pi N!} \left\| \int e^{-i(s,\omega)\cdot\xi} \left(\frac{\partial^2}{\partial s\partial\omega}\right)^N \left(b_j(s,x,(-\omega+\mu,p'),\hbar^{-1}\mu\ ;\ \hbar)\right) ds d\omega \right\|_{L^1_\xi},$$

where $\|\cdot\|_{L^1_\xi}$ denotes the L^1-norm of a function of the ξ variables. The well-known inequality

$$||\hat{v}||_{L^1(\mathbb{R}^d)} \le \sum_{|\alpha|\le d+1} ||\partial^\alpha v||_{L^1(\mathbb{R}^d)} \tag{4.6}$$

implies that the above bound is in turn bounded by

$$\frac{\hbar^N}{2\pi N!} \sum_{|\alpha|\le 3} \left\| \partial^\alpha \left(\frac{\partial^2}{\partial s\partial\omega}\right)^N \left[b_j(s,x,(-\omega+\mu,p'),\hbar^{-1}\mu\ ;\ \hbar)\right] \right\|_{L^1_{s,\omega}}$$

$$= \frac{\hbar^N}{2\pi N!} \sum_{|\alpha|\le 3} \Bigg\| \frac{(1-\rho(\mu))(-i\hbar^{-1})^{2K-j}}{(1+(\hbar^{-1}\mu)^2)^K} \partial^\alpha \left(\frac{\partial^2}{\partial s\partial\omega}\right)^N$$

$$\left[a_j(s,x,x,(-\omega+\mu,p'),\hbar^{-1}\mu)(-\omega+\mu)^{2K-j}\right] \Bigg\|_{L^1_{s,\omega}},$$

where $\|\cdot\|_{L^1_{s,\omega}}$ is defined similarly, and $\partial^\alpha = \frac{\partial^{\alpha_1}}{\partial s^{\alpha_1}} \frac{\partial^{\alpha_2}}{\partial \omega^{\alpha_2}}$, $\alpha = (\alpha_1, \alpha_2)$. Therefore B_j is bounded above by

$$\frac{\hbar^{-n+N-2K+j}}{(2\pi)^{n+1} N!} \sum_{|\alpha|\le 3} \int \left|\partial^\alpha \left(\frac{\partial^2}{\partial s \partial \omega}\right)^N \Big[\frac{(-\omega+\mu)^{2K-j} a_j(s,x,(-\omega+\mu,p'),\hbar^{-1}\mu)}{(1+(\hbar^{-1}\mu)^2)^k}\right.$$

$$\left.(1-\rho(\mu))\Big]\right| ds d\omega dp' d\mu$$

$$= \frac{\hbar^{-n+N+1-2K+j}}{(2\pi)^{n+1} N!} \sum_{|\alpha|\le 3} \int \left|\partial^\alpha \left(\frac{\partial^2}{\partial s \partial p_1}\right)^N \left[\frac{p_1^{2K-j} a_j(s,x,p,\sigma)}{(1+\sigma^2)^k}(1-\rho(\hbar\sigma))\right]\right| ds dp d\sigma.$$

Finally, notice that the integrand is $O\left(\sigma^{m'-2K}\right)$ and that $1-\rho(\hbar\sigma)$ has support in $|\sigma| \ge \hbar^{-1}\mu_0/2$. Therefore the above upper bound is less than a constant times

$$\begin{aligned} \hbar^{-n+N+1-2K+j} \int_{\hbar^{-1}\mu_0/2}^\infty \sigma^{m'-2K} d\sigma =& \hbar^{-n+N+1-2K+j} \frac{\sigma^{m'-2K+1}}{m'-2K+1} \Big|_{\hbar^{-1}\mu_0/2}^\infty \\ =& O\left(\hbar^{j-n-m'+N}\right). \end{aligned}$$

Since this is true for all positive integers N we are done. ☐

Lemma 4.3. *If $m' \ge 0$,*

(4.7)
$$\int u_\hbar(x,x)\,dx = \frac{1}{(2\pi\hbar)^n} \int 2\pi \; a(0,x,x,\mu,p',\hbar^{-1}\mu)\,dx\,dp'\,d\mu + O(\hbar^{-n-m'+1}).$$

Proof. By (4.4), it suffices to estimate

$$\tilde{u}_\hbar(x,x) = \frac{1}{(2\pi\hbar)^n} \int e^{-isp_1/\hbar + is\sigma} a(s,x,x,p,\sigma)\rho(\hbar\sigma) ds dp d\sigma.$$

By an argument identical to the one used in the proof of the Lemma,

$$\tilde{u}_\hbar(x,x) = \frac{1}{(2\pi\hbar)^n}\; \hbar^{-1} \int e^{is\omega/\hbar} a(s,x,x,(-\omega+\mu,p'),\hbar^{-1}\mu)\rho(\mu) ds dp' d\omega d\mu.$$

We will apply the method of stationary phase to the (s,ω) integral, before integrating with respect to p' and μ. To this end introduce the notation

$$u_\hbar(x,x,p',\mu) = \frac{1}{(2\pi\hbar)^n}\; \hbar^{-1} \int e^{is\omega/\hbar} a(s,x,x,(-\omega+\mu,p'),\hbar^{-1}\mu)\rho(\mu) ds d\omega,$$

so that the left-hand side of (4.7) is equal to $\int u_\hbar(x,x,p',\mu)\,dx dp' d\mu$ modulo $O(\hbar^\infty)$. We also have that

$$u_\hbar(x,x,p',\mu) = \frac{1}{(2\pi\hbar)^n} \int e^{-i\hbar\xi_1\xi_2} c(x,\mu,p';\; \hbar\;;\; \xi) d\xi,$$

where $\xi = (\xi_1, \xi_2)$ are the dual variables to (s, ω), and

$$c(x, \mu, p';\ \hbar\ ;\ \xi) = \frac{1}{2\pi} \int e^{-(s,\omega)\cdot\xi} a(s, x, x, (-\omega + \mu, p'), \hbar^{-1}\mu)\rho(\mu) ds d\omega$$

is the Fourier transform of ρa in (s, ω). Note that

$$\int c(x, \mu, p';\ \hbar;\ \xi) d\xi = 2\pi\ a(0, x, x, (\mu, p'), \hbar^{-1}\mu),$$

and therefore (by (4.5))

$$(4.8)\quad u_\hbar(x, x, p', \mu) - \frac{1}{(2\pi\hbar)^n}\ 2\pi\ a(0, x, x, (\mu, p'), \hbar^{-1}\mu) = \frac{1}{(2\pi\hbar)^n} \int R(\xi, \hbar) c(\xi) d\xi,$$

where $|R(\xi, \hbar)| \leq \hbar\, |\xi_1 \xi_2|$. Integrating (4.8) we see that the error term in (4.7) is bounded by

$$(4.9)\qquad \frac{1}{(2\pi\hbar)^n} \int_K \|Rc\|_{L^1_\xi}(x, p', \mu)\, dx\, dp'\, d\mu,$$

where K is a compact set containing the support of the left-hand side of (4.8). Using (4.6) again,

$$\begin{aligned} \hbar\, ||\xi_1 \xi_2 c||_{L^1_\xi} &= \frac{\hbar}{2\pi} \left|\left| \int e^{-i(s,\omega)\cdot\xi} \frac{\partial^2}{\partial s \partial \omega} a(s, x, x, (-\omega + \mu, p'), \hbar^{-1}\mu)\rho(\mu) ds d\omega \right|\right|_{L^1} \\ &\leq \frac{\hbar}{2\pi} \sum_{|\alpha| \leq 3} \left|\left| \partial^\alpha \frac{\partial^2}{\partial s \partial \omega} a(s, x, x, (-\omega + \mu, p'), \hbar^{-1}\mu)\rho(\mu) \right|\right|_{L^1_{(s,\omega)}}. \end{aligned}$$

Since $a(s, x, y, p, \sigma)$ is a classical symbol in σ of order m' and compactly supported in the rest of the variables, $\forall \alpha$ there exists a constant $C = C(\alpha)$ such that

$$(4.10)\qquad \begin{aligned} \left| \frac{\hbar}{2\pi} \partial^\alpha \frac{\partial^2}{\partial s \partial \omega} a(s, x, x, (-\omega + \mu, p'), \hbar^{-1}\mu)\rho(\mu) \right| &\leq C(\alpha) \left(1 + \hbar^{-1}|\mu|\right)^{m'} \hbar \\ &= C(\alpha) \left(\hbar + |\mu|\right)^{m'} \hbar^{-m'+1} \end{aligned}$$

for all (s, x, y, p). Integrating (4.10) with respect to (s, ω) over a sufficiently large compact set we obtain that

$$\|Rc\|_{L^1_\xi}(x, p', \mu) \leq C \left(\hbar + |\mu|\right)^{m'} \hbar^{-m'+1}.$$

We now integrate this inequality over the compact set, K, in (4.9), to obtain that the error term in (4.7) is bounded above by a constant times

$$(4.11)\qquad \hbar^{-m'-n+1} \int_{|\mu| \leq \mu_1} \left(\hbar + |\mu|\right)^{m'} d\mu$$

for some $\mu_1 > 0$ independent of $\hbar$, and this is $O(\hbar^{-m'-n+1})$ when $m' \geq 0$. □

Remark 4.4. Lemma 4.3 implies the Theorem (with a better error estimate in case $m' = 0$) if the amplitude a is homogeneous in the variable σ.

Proof of Theorem 4.1. Let us first assume that $m' > 0$. Since a is a symbol in σ of degree $m' > 0$, there exists $\tilde{a}(x,p)$ and a constant C such that

$$\left|a(0,x,x,p,\sigma) - \sigma^{m'}\tilde{a}(x,p)\chi(\sigma)\right| \le C\ (1+|\sigma|)^{m'-1},$$

where $\chi(\sigma)$ is smooth in $\sigma \neq 0$, and homogeneous of degree zero in σ. Then, in particular,

$$\left|\frac{2\pi}{(2\pi\hbar)^n}\ a(0,x,x,\mu,p',\hbar^{-1}\mu) - \frac{2\pi}{(2\pi\hbar)^n}\ \hbar^{-m'}\mu^{m'}\chi(\mu)\tilde{a}(0,x,(\mu,p'))\right|$$
$$\le (2\pi)^{1-n}C\ (1+\hbar^{-1}|\mu|)^{m'-1}\hbar^{-n}.$$

The left-hand side of this inequality is supported in $\mu \in [-\mu_0,\mu_0]$. After integrating with respect to μ, the remainder is bounded by constant times

$$\hbar^{-n}\int_{-\mu_0}^{\mu_0}(1+\hbar^{-1}|\mu|)^{m'-1}d\mu = 2\hbar^{-n-m'+1}\int_0^{\mu_0}(\hbar+\mu)^{m'-1}d\mu$$
$$= 2\hbar^{-n-m'+1}\left((\mu_0+\hbar)^{m'} - \hbar^{m'}\right)/m' = O(\hbar^{-n-m'+1}) \text{ since } m' > 0.$$

Therefore, for any ℓ' and $m' > 0$

$$\int u_\hbar(x,x)\,dx = (2\pi)^{-n}\hbar^{-\ell-m-n}\int 2\pi\ p_1^{m'}\tilde{a}(0,x,p)dxdp + O(\hbar^{-\ell-m-n+1}),$$

where $2\pi\ p_1^{m'}\tilde{a}(0,x,p)$ is the principal symbol on the diagonal.

Now let's assume $m' = 0$. Since a is a symbol in σ of degree zero, there exists $\tilde{a}(x,p)$ such that,

$$|a(0,x,x,p,\sigma) - \tilde{a}(x,p)\chi(\sigma)| < \frac{C}{1+|\sigma|}$$

for some constant $C > 0$.Then

$$\left|\frac{2\pi}{(2\pi\hbar)^n}a(0,x,x,\mu,p',\hbar^{-1}\mu) - \frac{2\pi}{(2\pi\hbar)^n}\tilde{a}(x,(\mu,p'))\chi(\mu)\right| < \frac{(2\pi)^{1-n}C}{1+\hbar^{-1}\,|\mu|}\ \hbar^{-n}$$

Again, since the left hand side is supported in the set $\{|\mu| \le \mu_0\}$, after integrating with respect to μ, the remainder is bounded by a constant times

$$\hbar^{-n}\int_0^{\mu_0}\frac{1}{1+\hbar^{-1}\mu}d\mu = \hbar^{-n+1}\int_0^{\mu_0}\frac{1}{\mu+\hbar}d\mu$$
$$= \hbar^{-n+1}\left(log(\mu_0+\hbar) - log(\hbar)\right) = O(\hbar^{-n+1}\ log(1/\hbar)).$$

Therefore, for $m = 1/2$

$$\int u_\hbar(x,x)\,dx = (2\pi)^{-n}\hbar^{-\ell-m-n}\int 2\pi\ \tilde{a}(0,x,p)dpdx + O(\hbar^{-\ell-m-n+1}\ \log(1/\hbar)),$$

where $2\pi\ \tilde{a}(0,x,p)$ is the principal symbol of the family on the diagonal.

4.2. **The trace in case $m' \leq -4$.**

Theorem 4.5. *With the previous notation, if $m \leq -7/2$, and $\widehat{A}$ an operator in $\mathcal{A}_{X_c}$, then each classical ΨDO $\sigma_1(A)_s$ obtained from the symbol of $\widehat{A}$ in the flow out on each orbit is of trace class, and*

$$\mathrm{Tr}(\widehat{A}) = (2\pi)^{-n+1/2}\hbar^{-n-\ell+1/2}\int_S \mathrm{Tr}\,(\sigma_1(A)_s)\,ds + O(\hbar^{-n-\ell+3/2}). \tag{4.12}$$

Lemma 4.6. *If $m' < 0$ and $\ell' = 0$,*

$$\int u_\hbar(x,x)\,dx = \frac{1}{(2\pi\hbar)^n}2\pi\int a(0,x,x,\mu,p',\hbar^{-1}\mu)\,dx\,dp'\,d\mu + O(\hbar^{-n+2}). \tag{4.13}$$

Proof. This follows immediately from (4.11) (which was derived under no assumptions on m'). ☐

Proof. Starting with equation (4.3), since $m' \leq -4$

$$a_2(s,x,y,p) := \int e^{is\sigma}a(s,x,y,p,\sigma)d\sigma$$

is absolutely convergent and can be extended to a C^2 function of s. In addition, a_2 is compactly supported in s,x,y,p. Using the stationary phase theorem for C^{2k} amplitudes (Theorem 7.7.5 in [18]), we get

$$\int e^{i\hbar^{-1}sp_1}a_2(s,x,x,-p_1,p')dsdp_1 \sim 2\pi\,\hbar\,a_2(0,x,x,0,p'),$$

and then

$$\int u_\hbar(x,x)\,dx = (2\pi)^{-n+1/2}\hbar^{-n-\ell+1/2}\int\sqrt{2\pi}\;a_2(0,x,x,0,p')dxdp' + O(\hbar^{-n-\ell+3/2}),$$

where $\sqrt{2\pi}\;a_2(0,x,x,0,p') = \sqrt{2\pi}\int e^{is\sigma}a(s,x,x,p)d\sigma\Big|_{s=0,p_1=0}$ is the extension of the symbol σ_1 to the intersection of the Lagrangians. This is the desired result in the model case. ☐

5. PROJECTORS AND "CUT" QUANTUM OBSERVABLES

In this section we will prove that, under a mild additional condition on ∂X_c, the algebra $\mathcal{A}_{X_c}$ contains orthogonal projectors. We will also prove that, in case there exists an $\hbar$-pseudodifferential operator on M, $\hat{P}$, such that:

1. The spectrum of $\hat{P}$ is discrete and is contained in $\hbar\mathbb{Z}$, and
2. $X_c = P^{-1}(I)$ for $I \subset \mathbb{R}$ a closed interval,

then the spectral projector of $\hat{P}$ associated to the interval I is in $\mathcal{A}_{X_c}$

5.1. **On the Existence of Projectors.** In addition to the assumptions on ∂X_c that we have been making throughout, let us now assume that ∂X_c is of contact type. Recall that this means that there exists a one-form β on ∂X_c such that (a) $d\beta$ is the pull-back of the symplectic form to ∂X_c, and (b) $\beta \rfloor \Xi_P$ is constant, where P is a defining function of X_c with periodic flow on ∂X_c.

Following the proof of Lemma 5 in [4], one obtains:

Lemma 5.1. *There exists an smooth function, which will be called again* $P : T^*M \to \mathbb{R}$, *such that*

- (a) P *is bounded from below and tends to* ∞ *at infinity in the cotangent directions,*
- (b) $\partial X_c = P^{-1}(0)$, *and*
- (c) *There exists a neighborhood* $\mathcal{W}$ *of* ∂X_c *such that the Hamilton flow of* P *is* 2π*-periodic in* $\mathcal{W}$.

Next we recall (see [17] Proposition 3.8) how to obtain a quantum version of the previous result:

Lemma 5.2. *Let* $\widehat{P}(\hbar)$ *be a semiclassical pseudodifferential operator with principal symbol* P *and vanishing sub-principal symbol. Let* μ *be the Maslov index of the trajectories of* Φ^P *(the Hamilton flow of* P*) in* $\mathcal{W}$. *Assume the Bohr-Summerfeld conditions* (3.1). *Then there exists a semiclassical pseudodifferential operator* $\widehat{R}_2(\hbar)$ *of order* -2 *such that for* $\epsilon << 1$

$$Spec\left(P - \frac{\mu}{4}\,\hbar - \widehat{R}_2(\hbar)\right) \cap [-\epsilon/3, \epsilon/3] \subset \hbar\mathbb{Z}$$

when we restrict $\hbar$ *to the sequence* $\hbar = 1/N$ *with* N *large.*

Proof. Pick $\epsilon > 0$ such that $P^{-1}[-\epsilon, \epsilon] \subset \mathcal{W}$. Let ρ be a smooth function with support in $[-\epsilon, \epsilon]$, such that $\rho \equiv 1$ on $[-\epsilon/2, \epsilon/2]$. Let

$$\gamma = \frac{1}{2\pi} \int_0^{2\pi} p(t)\dot{x}(t) - P(x(t), p(t))dt$$

be the (common) action of the trajectories of the Hamilton flow of P in $\mathcal{W}$. Then $e^{-2\pi i \hbar^{-1}\left(\widehat{P} - \frac{\mu}{4}\,\hbar - \gamma\right)}$ is microlocally in $\mathcal{W}$ a pseudodifferential operator with symbol identically equal to one, and thus one can write

$$\rho(\widehat{P}) e^{-2\pi i \hbar^{-1}\left(\widehat{P} - \frac{\mu}{4}\,\hbar - \gamma\right)} = \rho(\widehat{P})(I + \hbar \widehat{R}(\hbar)),$$

where $\widehat{R}(\hbar)$ is a zeroth order $\hbar$-ΨDO. Recall that γ is an integer in ∂X_c, and therefore, for $\hbar = 1/N$ we obtain

$$\rho(\widehat{P}) e^{-2\pi i \hbar^{-1}\left(\widehat{P} - \frac{\mu}{4}\,\hbar\right)} = \rho(\widehat{P})(I + \hbar \widehat{R}(\hbar)),$$

Since $I + \hbar\widehat{R}(\hbar)\rho(\widehat{P})$ has spectrum close to 1 for $\hbar << 1$, one can then define for $\hbar$ small

$$\widehat{R}_2 = -\frac{\hbar}{2\pi i}\log(I + \hbar\widehat{R}(\hbar)\rho(\widehat{P})),$$

and since $\widehat{R}_2$ commutes with $\widehat{P}$, we obtain

$$\rho(\widehat{P})e^{-2\pi i\hbar^{-1}(\widehat{P}-\frac{\mu}{4}\ \hbar-\widehat{R}_2)} = \rho(\widehat{P})\left(I + \hbar\widehat{R}(\hbar)\right)\left(I + \hbar\widehat{R}(\hbar)\rho(\widehat{P})\right)^{-1}.$$

Since $\rho \equiv 1$ on $[-\epsilon/2, \epsilon/2]$, the spectral theorem guarantees that the above operator is the identity on any eigenfunction of $\hat{P}$ with eigenvalue in $[-\epsilon/2, \epsilon/2]$. Since, for $\hbar$ small enough, the spectrum of $\widehat{P} - \frac{\mu}{4}\ \hbar - \widehat{R}_2$ in $[-\epsilon/3, \epsilon/3]$ corresponds to eigenfunctions of $\hat{P}$ with eigenvalues in $[-\epsilon/2, \epsilon/2]$, the result follows. ☐

Let us define $\widehat{P}_2 := \widehat{P} - \frac{\mu}{4}\ \hbar - \widehat{R}_2$, whose principal symbol continues to be P. Let χ be the characteristic function on $(-\infty, 0]$, and define the projector

$$\Pi = \chi\left(\widehat{P}_2\right)$$

Theorem 5.3. *The projector* Π *defined above belongs to the class* $J^{-1/2,1/2}(M \times M, \Delta, \mathcal{F}\partial X_c)$.

Proof. Let ρ be the cut-off function of the previous proof. We decompose

$$\Pi = (1-\rho)\chi\left(\widehat{P}_2\right) + \rho\chi\left(\widehat{P}_2\right)$$

Clearly $(1-\rho)\chi\left(\widehat{P}_2\right)$ is a semiclassical pseudodifferential operator. Therefore we need to prove that $\rho\,\chi\left(\widehat{P}_2\right)$ belongs to $J^{-1/2,1/2}$. This operator has microsupport in $\mathcal{W}$.

We take $\mathbb{R}^{n-1} \times S^1$ as the model case with coordinates (x, θ) and $T^*(\mathbb{R}^{n-1} \times S^1)$ with coordinates $(x, \theta; p, \tau)$. Let $\Pi^E = \Pi^E_N$ be the projector on eigenfunctions of $\widehat{P}^n := \hbar D_\theta = \frac{\hbar}{i}\frac{\partial}{\partial\theta}$ with eigenvalues greater than or equal to E/N, where $E \in \mathbb{Z}$ is a constant. (Here θ is the 2π-periodic variable in S^1.) Let $T_s = e^{-is\hbar^{-1}\widehat{P}_2}$, and let

$$T^n_s = e^{-i\hbar^{-1}s\ \widehat{P}^n}$$

be the translation representation on $L^2(\mathbb{R}^{n-1} \times S^1)$. Let $(x_0, p_0) \in \mathcal{W}$. As in [13, 7], there exist an S^1-invariant neighborhood $\mathcal{U} \subset \mathcal{W}$ of (x_0, p_0) (the circle action given by the Hamilton flow of P), an S^1-invariant open set $\mathcal{U}^n \subset T^*(\mathbb{R}^{n-1} \times S^1)$, and an S^1-equivariant canonical transformation

$$\phi : \mathcal{U} \to \mathcal{U}^n,$$

which sends $\partial X_c \cap \mathcal{U}$ into $\left\{(x,\theta;p,\tau) \in T^*\left(\mathbb{R}^{n-1}\times S^1\right) \;\middle|\; \tau = E\right\}$. Again, as in [13, 7], one can show that there exists a semiclassical zeroth order Fourier integral operator

$$F : L^2(M) \to L^2(\mathbb{R}^{n-1}\times S^1),$$

with microsupport on $\mathcal{U}\times\mathcal{U}^n$ such that

$$F^*F = I_{\mathcal{U}^n},\ FF^* = I_{\mathcal{U}},$$

and

$$F\rho\, \chi(\widehat{P}_2) = \rho\left(\Pi^E\right)F,$$

This reduces the proof to the model case. It suffices to show that $\Pi\widehat{Q}$ is on the algebra, for any zeroth order compactly supported semiclassical pseudodifferential operator $\widehat{Q}$ in $\mathbb{R}^{n-1}\times S^1$. Note that

$$\Pi^E\ \widehat{Q} = \frac{1}{2\pi}\int_0^{2\pi} e^{-iNs\widehat{P}^n} e^{iNsE}\ \widehat{Q}\ \frac{1}{1-e^{is}}ds$$

The operator $e^{-iNs\widehat{P}^n}e^{iNsE}\widehat{Q}$ is a semiclassical FIO with Lagrangian

$$\left\{((x,\theta\ ;\ p,\tau=E),(y=x,\alpha+s=\theta\ ;\ -p,-\tau=-E))\,\middle|\,x\in\mathbb{R}^{n-1},\theta,\alpha\in[0,2\pi]\right\}.$$

Therefore, in local coordinates, the Schwartz kernel of $\Pi^E\widehat{Q}$ can be written as

$$\frac{1}{(2\pi\hbar)^n}\frac{1}{2\pi}\int e^{iN(\ (x-y)p+(\theta-\alpha-s)(\tau-E)\ )}q(x,\theta,y,\alpha,p,\tau,\hbar)\frac{1}{1-e^{is}}dpd\tau ds, \tag{5.1}$$

where q is symbol with expansion in $\hbar$. Notice that $\frac{1}{1-e^{is}}$ is a conormal distribution in $s=0$ and equation (5.1) shows the hybrid nature of the amplitude of the projector. Equation (5.1) also proves that $\Pi^E\widehat{Q}\in J^{-1/2,1/2}$, with principal symbol $\chi_{X_c}Q(x,\theta\ ;\ p,\tau)$ in the diagonal, and

$$\frac{1}{\sqrt{2\pi}}\frac{Q(\Phi_s^P(x,\theta\ ;\ p,\tau))}{1-e^{is}}$$

in the flow-out. □

Remark 5.4. If one has an $\hbar$-pseudodifferential operator $\widehat{P}$ with discrete spectrum such that

$$\mathrm{Spec}(\widehat{P}(\hbar))\subset\hbar\mathbb{Z},$$

then the previous proof shows that, for any given $E_1,E_2\in\mathbb{Z}$ such that $E_1<E_2$, and for $j=1,2$:

> the trajectories on $P^{-1}(E_j)$ satisfy the Bohr-Sommerfeld condition (3.1).

Then the orthogonal projector onto

$$\mathcal{H}_N = \text{span of eigenvectors of } \widehat{P}(\hbar) \text{ with eigenvalues in } [E_1,E_2],$$

is in the algebra $J^{-1/2,1/2}\left(M \times M; \Delta, \mathcal{F}\partial X_c\right)$, associated to

$$X_c = \left\{ \overline{x} \in T^*M \;\middle|\; E_1 \le P(\overline{x}) \le E_2 \right\}.$$

5.2. **Cut quantum observables.** In this section we fix a projector Π as in §5.1, and consider "cut" quantum observables, by which we mean operators of the form

$$\Pi \widehat{Q} \Pi$$

where Q is a pseudodifferential operator on M. By Theorem 5.3 these operators are in $J^{-1/2,1/2}\left(M \times M; \Delta, \mathcal{F}\partial X_c\right)$. The symbolic properties of these operators are summarized by the following Proposition:

Proposition 5.5. *Let $\widehat{Q}(\hbar)$ be a zeroth-order semiclassical pseudodifferential operator with compact microsupport. Then $\Pi\widehat{Q}\Pi$ is in the class $J^{-1/2,1/2}(M \times M \,;\, \Delta, \mathcal{F}\partial X_c)$. Its symbols, ignoring Maslov factors, are as follows:*

$$\sigma_0(\Pi\widehat{Q}\Pi)(\overline{x},\overline{x}) = \chi_{X_c}(\overline{x}) Q(\overline{x}) \, \sqrt{dx \wedge dp}, \tag{5.2}$$

$$\sigma_1\left(\Pi\widehat{Q}\Pi\right)_s = \Pi_{F_s} M_{|_{Q_{F_s}}} \Pi_{F_s} \tag{5.3}$$

where χ_{X_c} is the characteristic function of X_c, $\overline{x} = (x,p) \in T^*M$, F_s is the fiber above $s \in S$, $Q_{|_{F_s}}$ is the restriction of Q to F_s, $M_{Q_{|_{F_s}}}$ is the operator "multiplication by $Q_{|_{F_s}}$", and Π_{F_s} is the Szegö projector in the orbit F_s, i.e., for $u : F_s \to \mathbb{C}$ smooth, $u(\Phi^P_s \bar{y}) = \sum_j u_j(\bar{y}) \frac{e^{ijs}}{\sqrt{2\pi}}$, $[\Pi_{F_s} u](\bar{x}) = \sum_{j \ge 0} \frac{u_r(\overline{x})}{\sqrt{2\pi}}$.

Remark 5.6. Notice that the Szegö projector Π_{F_s} is a classical pseudodifferential operator with principal symbol $\chi_{(T^*F_s)^+}$, where $(T^*F_s)^+$ is the part with positive momentum variable, in the direction of the Hamilton flow. This function is smooth in $T^*F_s \setminus 0$.

Remark 5.7. The symbol of $\Pi_{|_{F_s}} M_{Q_{|_{F_s}}} \Pi_{|_{F_s}}$ is $\chi_{(T^*F_s)^+} Q_{|_{F_s}}$, which agrees with the symbol in the diagonal, restricted to the intersection. This is the so-called symbolic compatibility condition referred to in Theorem 3.17.

Proof. The first part was proven in Theorem 5.3. The principal symbol in the diagonal is clear. Using the relation (0.2) in [1], we obtain that for $\overline{y} \in \partial X_c$, $s \neq 0$,

$$\begin{aligned}\sigma_1(\Phi^P_s(\overline{y}) \,;\, \overline{y}) &= \frac{1}{\sqrt{2\pi}} \frac{1}{2\pi} \int_0^{2\pi} \frac{Q(\Phi^P_{s-\tilde{s}}(\overline{y}))}{1 - e^{i(s-\tilde{s})}} \frac{1}{1 - e^{i\tilde{s}}} d\tilde{s} \\ &= \frac{1}{\sqrt{2\pi}} \frac{1}{2\pi} \int_0^{2\pi} \frac{Q(\Phi^P_{\tilde{s}}(\overline{y}))}{1 - e^{i\tilde{s}}} \, \frac{1}{1 - e^{i(s-\tilde{s})}} d\tilde{s}.\end{aligned}$$

Since $Q(\Phi^P_{\tilde{s}}(\overline{y}))$ is a smooth 2π-periodic function in $\tilde{s}$, there exists a sequence of functions $\{Q_j(\overline{y})\}_{j=-\infty}^{\infty}$ such that

$$Q(\Phi^P_{\tilde{s}}(\overline{y})) = \sum_{j=-\infty}^{\infty} Q_j(\overline{y}) \frac{e^{ij\tilde{s}}}{\sqrt{2\pi}},$$

and the symbol becomes:

$$\begin{aligned}
\sigma_1(\Phi^P_s(\overline{y}) \ ; \ \overline{y}) &= \frac{1}{\sqrt{2\pi}} \frac{1}{2\pi} \int \left(\sum_j \frac{Q_j(\overline{y})}{\sqrt{2\pi}} e^{ij\tilde{s}} \right) \left(\sum_{k \geq 0} e^{ik\tilde{s}} \right) \left(\sum_{k' \geq 0} e^{ik'(s-\tilde{s})} \right) d\tilde{s} \\
&= \frac{1}{\sqrt{2\pi}} \frac{1}{1-e^{is}} \left[\sum_{j \geq 0} e^{ijs} \frac{Q_j(\overline{y})}{\sqrt{2\pi}} + \sum_{j<0} \frac{Q_j(\overline{y})}{\sqrt{2\pi}} \right].
\end{aligned}$$

We now interpret this as the kernel of an operator acting on the fibers of $\partial X_c \to S$. Let us consider a fiber $F_s \subset \partial X_c$, and a function $u : F_s \to \mathbb{C}$. For every fixed $\overline{y} \in \partial X_c$, let $u_r(\overline{y})$ and $Q_j(\overline{y})$ be the corresponding Fourier coefficients in each decomposition

$$u(\Phi^P_s \overline{y}) = \sum_{r=-\infty}^{\infty} u_r(\overline{y}) \frac{e^{irs}}{\sqrt{2\pi}}, \ Q(\Phi^P_s \overline{y}) = \sum_{j=-\infty}^{\infty} Q_j(\overline{y}) \frac{e^{ijs}}{\sqrt{2\pi}}$$

As a pseudodifferential operator, the symbol in in the flow-out of $\Pi\widehat{Q}\Pi$ acting the function u, is given by

(5.4)

$$\begin{aligned}
&\left[\sigma_1 \left(\Pi\widehat{Q}\Pi\right)_s \ u\right](\overline{x}) = \int_{F_s} \sigma_1(\Pi\widehat{Q}\Pi)(\overline{x},\overline{y}) u(\overline{y}) d\overline{y} = \int_{S^1} \sigma_1(\Pi\widehat{Q}\Pi)(\overline{x}, \Phi^P_s \overline{x}) u(\Phi^P_s \overline{x}) ds \\
&= \int \frac{1}{\sqrt{2\pi}} \frac{1}{1-e^{-is}} \left[\sum_{j \geq 0} \frac{Q_j(\overline{x})}{\sqrt{2\pi}} + \sum_{j<0} \frac{Q_j(\overline{x})}{\sqrt{2\pi}} e^{ijs} \right] \sum_r u_r(\overline{x}) \frac{e^{irs}}{\sqrt{2\pi}} ds \\
&= \sum_{r \geq 0, j \geq -r} u_r(\overline{x}) \frac{Q_j(\overline{x})}{\sqrt{2\pi}}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&\left[\Pi_{F_s} M_{|_{Q_{F_s}}} \Pi_{F_s} \ (u)\right](\overline{x}) = \Pi_{F_s} M_{|_{Q_{F_s}}} \left(\sum_{r \geq 0} \frac{u_r(\overline{y})}{\sqrt{2\pi}} \right)(\overline{x}) = \Pi_{F_s} \left(Q(\overline{y}) \sum_{r \geq 0} \frac{u_r(\overline{y})}{\sqrt{2\pi}} \right)(\overline{x}) \\
&\quad = \Pi_{S^1} \left(Q(\Phi^P_s \overline{x}) \sum_{r \geq 0} \frac{u_r(\Phi^P_s \overline{x})}{\sqrt{2\pi}} \right)_{\Big|_{s=0}} = \Pi_{S^1} \left(\sum_{j \in \mathbb{Z}, r \geq 0} \frac{Q_j(\overline{x})}{\sqrt{2\pi}} u_r(\overline{x}) \frac{e^{i(j+r)s}}{\sqrt{2\pi}} \right)_{\Big|_{s=0}}
\end{aligned}$$

$$= \sum_{r\geq 0, j\geq -r} Q_j(\overline{x}) \frac{u_r(\overline{x})}{\sqrt{2\pi}},$$

which agrees with equation (5.4). This proves

$$\sigma_1\left(\Pi\widehat{Q}\Pi\right)_s = \Pi_{F_s} M_{|_{Q_{F_s}}} \Pi_{F_s},$$

which yields (5.3) after applying Proposition 3.18.

□

Operators of the form $\Pi_N \widehat{Q}_N \Pi_N : \mathcal{H}_N \to \mathcal{H}_N$ generalize Toeplitz matrices, and this is reflected in its principal symbol in $\mathcal{F}\partial X_c$. (In case $M = S^1$ and Π_N the projector onto the span of $\{e^{ij\theta},\ j = 0, \dots, N\}$, the $\Pi_N \widehat{Q}_N \Pi_N$ are to leading order the generalized Toeplitz matrices of [6], page 84.)

5.2.1. *Applications: A symbolic proof of the Szegö Limit Theorem.* We begin with the functional calculus, the heart of which is the following

Lemma 5.8. *Let $\widehat{Q}$ be a self-adjoint pseudodifferential operator of order zero on M. Then*

$$\Pi\, e^{-it\Pi\widehat{Q}\Pi} \in J^{-1/2,1/2}\left(M \times M; \Delta, \mathcal{F}\partial X_c\right).$$

Proof. Let us define

$$W(t) = \Pi e^{-it\Pi\widehat{Q}\Pi}.$$

It is the solution of the problem

$$\text{(5.5)} \qquad \begin{cases} \frac{1}{i}\frac{\partial}{\partial t} W(t) + \Pi\widehat{Q}\Pi W(t) = 0 \\ W(t)\Big|_{t=0} = \Pi \end{cases}$$

The idea in the following proof is to construct a solution which will be in the algebra and will make the right-hand side of first equation (5.5) of order $O(\hbar^\infty)$.

As a first approximation we take

$$\widetilde{W_0} = \Pi e^{-it\widehat{Q}},$$

which satisfies

$$\frac{1}{i}\frac{\partial}{\partial t}\widetilde{W_0} + \Pi\widehat{Q}\Pi\, \widetilde{W_0} = -\Pi\left[\Pi, \widehat{Q}\right] e^{-it\widehat{Q}}.$$

We will prove below that $\left[\Pi, \widehat{Q}\right] \in \text{sc-}I^{-1/2}(M \times M; \mathcal{F}\partial X_c)$ (see Section 5.2.2). Therefore, we obtain

$$\begin{cases} \frac{1}{i}\frac{\partial}{\partial t}\widetilde{W_0}(t) + \Pi\widehat{Q}\Pi\widetilde{W_0}(t) =: \widetilde{R}_0(t) \in \text{sc-}I^{-1/2}\left(M \times M; \mathcal{F}\partial X_c\right) \\ \widetilde{W_0}(t)\Big|_{t=0} = \Pi \end{cases}$$

We will now modify W_0 so as to make the right hand side $O(\hbar^\infty)$ instead of an operator in sc-$I^{-1/2}(M\times M;\mathcal{F}\partial X_c)$. For the rest of the proof, it will be convenient to identify symbols in the flow-out with corresponding families of smoothing operators acting on functions on the fibers F_s for each $s\in S$. The symbol $R_0(t)$ has a corresponding family operator $\{\widetilde{\mathcal{R}}_{0,s}(t)\}_{s\in S}$. Let us consider the following problem,

$$\begin{cases} \frac{1}{i}\frac{\partial}{\partial t}\widetilde{\mathcal{V}}_{0,s}+\Pi_{F_s}M_{Q_{|_{F_s}}}\Pi_{F_s}\circ\widetilde{\mathcal{V}}_{0,s}=-\widetilde{\mathcal{R}}_{0,s}, \\ \\ \widetilde{\mathcal{V}}_{0,s}\Big|_{t=0}=0, \end{cases}$$

whose solution is

$$\widetilde{\mathcal{V}}_{0,s}(t)=-i\int_0^t e^{i(\tilde{t}-t)\Pi_{F_s}M_{Q_{|_{F_s}}}\Pi_{F_s}}\widetilde{\mathcal{R}}_{0,s}(\tilde{t})d\tilde{t}.$$

Notice that $\widetilde{\mathcal{V}}_{0,s}$ is a smoothing operator. Let us call V_0 the smooth symbol in $\mathcal{F}\partial X_c$ given by the operator $\widetilde{\mathcal{V}}_{0,s}$. Take $\widetilde{V_0}\in$ sc-$I^{-1/2}(M\times M;\mathcal{F}\partial X_c)$ with symbol V_0. By construction

$$\frac{1}{i}\frac{\partial}{\partial t}\left(\widetilde{W_0}+\widetilde{V_0}\right)+\Pi\widehat{Q}\Pi\left(\widetilde{W_0}+\widetilde{V_0}\right)=\hbar\widetilde{R}_1(t)\in \text{sc-}I^{-3/2}\left(M\times M;\mathcal{F}\partial X_c\right).$$

Proceeding inductively one can find a sequence of operators $\tilde{V}_j$ such that for all J

$$\frac{1}{i}\frac{\partial}{\partial t}\left(\widetilde{W_0}+\sum_{j=0}^{J}\hbar^j\ \widetilde{V_j}\right)+\Pi\widehat{Q}\Pi\left(\widetilde{W_0}+\sum_{j=0}^{J}\hbar^j\ \widetilde{V_j}\right) \tag{5.6}$$
$$=\hbar^{J+1}\widetilde{R}_{J+1}\in\text{sc-}I^{-3/2-J}\left(M\times M;\mathcal{F}\partial X_c\right)$$

Finally, take an operator $\widetilde{V}\in$ sc-$I^{-1/2}(M\times M;\mathcal{F}\partial X_c)$ such that $\widetilde{V}\sim\sum_{j=0}^{\infty}\widetilde{V}_j$, and define $\widetilde{W}=\widetilde{W_0}+\widetilde{V}$. Then

$$\frac{1}{i}\frac{\partial}{\partial t}\widetilde{W}+\Pi\widehat{Q}\Pi\widetilde{W}=O(\hbar^\infty).$$

A standard application of Duhamel's principle finishes the proof. □

Proposition 5.9. *Let $\widehat{Q}$ be a self-adjoint semiclassical pseudodifferential operator. Then for any smooth function f, $\Pi\, f(\Pi\widehat{Q}\Pi)$, is in the class $J^{-1/2,1/2}(M\times M\ ;\ \Delta,\mathcal{F}\partial X_c)$. The symbols, ignoring Maslov factors, are as follows:*

$$\sigma_0\left(\Pi\, f(\Pi\widehat{Q}\Pi)\right)(\overline{x},\overline{x})=\chi_{X_c}(\overline{x})f\left(Q(\overline{x})\right)\ \sqrt{dx\wedge dp},\ \text{and} \tag{5.7}$$

$$\sigma_1\left(\Pi\, f(\Pi\widehat{Q}\Pi)\right)\Big|_{F_s}=\Pi_{F_s}\, f\left(\Pi_{F_s}M_{Q_{|_{F_s}}}\Pi_{F_s}\right), \tag{5.8}$$

where F_s,$Q_{|_{F_s}}$, $M_{Q_{|_{F_s}}}$, and Π_{F_s} are as in Proposition 5.5

Proof. We have:

$$\Pi\, f\left(\Pi\widehat{Q}\Pi\right) = \frac{1}{\sqrt{2\pi}} \int \Pi\, e^{-it\Pi\widehat{Q}\Pi} \check{f}(t)dt, \tag{5.9}$$

where $\check{f}(t) = \frac{1}{\sqrt{2\pi}}\int e^{is} f(s)ds$. By the previous lemma we can conclude that $\Pi f(\Pi\widehat{Q}\Pi) \in J^{-1/2,1/2}$. Moreover

$$\sigma_0(\Pi f(\Pi\widehat{Q}\Pi))(\overline{x},\overline{x}) = \frac{1}{\sqrt{2\pi}} \int e^{-itQ(\overline{x})} \chi_{X_c}(\overline{x})\check{f}(t)dt = f(Q(\overline{x}))\ \chi_{X_c}(\overline{x}),$$

and

$$\sigma_1\left(\Pi f(\Pi\widehat{Q}\Pi)\right)_s = \frac{1}{\sqrt{2\pi}} \int \Pi_{F_s} e^{-it\Pi_{F_s}M_{Q_{F_s}}\Pi_{F_s}} \check{f}(t)dt = \Pi_{F_s} f(\Pi_{F_s}M_{Q_{F_s}}\Pi_{F_s}).$$

□

As an immediate corollary of Theorems 4.1 and 5.5, we obtain the following Szegö limit theorem:

Corollary 5.10. *Assume that X_c is compact. Then for any smooth function f*

$$\operatorname{Tr}\left(\Pi_N f(\Pi_N\widehat{Q}_N\Pi_N)\right) = (2\pi)^{-n}\ N^n \int_{X_c} f\circ Q\ \frac{\omega^n}{n!} + O(N^{n-1}\log(N)).$$

5.2.2. *Commutators.* We now describe another property of the projector. Let $\widehat{Q}$ be a semiclassical pseudodifferential operator as above. The projector Π behaves microlocally as the identity on the interior X_c, suggesting that $\left[\Pi,\widehat{Q}\right]$ is microlocally $O(\hbar^\infty)$ on the diagonal. Using Proposition 3.14, we anticipate that $\left[\Pi,\widehat{Q}\right] \in \text{sc-}I^{-1/2}\left(M\times M;\mathcal{F}\partial X_c\right)$. We now prove that this is indeed the case, and compute the principal symbol of the commutator.

Proposition 5.11. *For any zeroth order compactly supported semiclassical pseudodifferential operator $\widehat{Q}$, $\left[\Pi,\widehat{Q}\right] \in$ sc-$I^{-1/2}\left(M\times M;\mathcal{F}\partial X_c\right)$ is a semiclassical Fourier integral operator, with (smooth) principal symbol*

$$\sigma\left(\left[\Pi,\widehat{Q}\right]\right)\left(\overline{x}=\Phi^P_s\overline{y}\ ;\ \overline{y}\right) = \begin{cases} \frac{1}{\sqrt{2\pi}}\frac{Q(\overline{y})-Q(\overline{x})}{1-e^{is}} & \text{if } \overline{x}\neq\overline{y} \\ \\ \frac{1}{i\sqrt{2\pi}}\left\{P,Q\right\}(\overline{x}) & \text{if } \overline{x}=\overline{y}. \end{cases} \tag{5.10}$$

Furthermore, if the principal symbol of $\widehat{Q}$ is constant along the orbits in the flow-out, the Hamilton flows Φ^P and Φ^Q of P and Q commute in ∂X_c, and the subprincipal symbol of Q vanishes on ∂X_c (Levi condition), then $\left[\Pi,\widehat{Q}\right] \in I^{-5/2}\left(M\times M;\mathcal{F}\partial X_c\right)$.

Remark 5.12. The Hamilton flow of the principal symbol of an operator that commutes with Π preserves the region X_c. However, the converse is not true; the invariance of the region X_c is a much weaker condition than the commuting property.

Proof. It is enough to prove it in the model case $\mathbb{R}^{n-1} \times S^1$ with coordinates (x, θ) and $T^*(\mathbb{R}^{n-1} \times S^1)$ with coordinates $(x, \theta\,; p, \tau)$. We only consider one energy level, say E so that $X_c = \{\tau \geq E\}$. For simplicity, let us consider zeroth order semiclassical pseudodifferential operators of the form:

$$\widehat{Q}(\hbar) f(x, \theta) = \sum_m e^{im\theta} \int q(x, p, \theta, \hbar m) e^{ixp/\hbar} \hat{f}(p, m, \hbar) dp, \text{ where}$$

$$\hat{f}(p, m, \hbar) = \frac{1}{(2\pi\hbar)^{n+1}} \int e^{-iyp/\hbar} e^{-im\alpha} f(y, \alpha) dy d\alpha,$$

and $q(x, p, \theta, s)$ is the full symbol. Decompose q in its Fourier modes, $q_k(x, p, \theta, s) = \sum e^{ik\theta} q_k(x, p, s)$. Then $\widehat{Q} = \sum \widehat{Q}_k$, where

$$\widehat{Q}_k(\hbar) f = \sum_m e^{im\theta} \int e^{ik\theta} q_k(x, p, \hbar m) e^{ixp/\hbar} \hat{f}(p, m, \hbar) dp \tag{5.11}$$

is a sc-ΨDO with symbol $q_k(x, p, s)$. The kernel of Q_k is

$$K_{Q_k}(x, y, \theta, \alpha) = \sum_m e^{i(k+m)\theta - im\alpha} p_k(x, y, m, \hbar),$$

where

$$p_k(x, y, m, \hbar) = \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i(x-y)p/\hbar} q_k(x, p, \hbar m) dp.$$

A calculation shows that for $k > 0$,

$$K_{[\Pi_N, \widehat{Q}_k(1/N)]}(x, \theta, y, \alpha)$$

$$= \frac{N^{n+1}}{(2\pi)^{n+1}} \int e^{iN(x-y)p} e^{iNE(\theta-\alpha)} \left[\sum_{0<j\leq k} e^{-ij(\theta-\alpha)} e^{ik\theta} \; q_k(x, p, E - j/N) \right] dp.$$

Notice that the amplitude in the integral above (in brackets) has an expansion in powers of $\hbar$, and in fact is a semiclassical symbol. The phase parametrizes the flow-out of $\{\tau = E\}$ by the canonical S^1 action, which is $\mathcal{F}\partial X_c$. This proves that the commutator is in the corresponding class, and the principal symbol is

$$\frac{1}{\sqrt{2\pi}} \sum_{0<j\leq k} e^{-ij(\theta-\alpha)} e^{ik\theta} q_k(x, p, E) = \frac{q_k(x, p, E) e^{ik\alpha} - q_k(x, p, E) e^{ik\theta}}{1 - e^{-i(\alpha-\theta)}}.$$

The case $k < 0$ is similar, and taking the sum over k, we obtain (5.10).

For the last part, assume that the principal symbol is constant in the fibers of $\partial X_c \to S$, which implies that $\left[\Pi, \widehat{Q}\right] \in \text{sc-}I^{-3/2}\left(M \times M; \mathcal{F}\partial X_c\right)$. Assuming the Levi condition, we will show next that the principal symbol (corresponding to the degree $-3/2$) vanishes again. Take $\overline{x}_0, \overline{y}_0 \in \mathcal{F}\partial X_c$, $\overline{x}_0 \neq \overline{y}_0$. Consider two zeroth order semiclassical pseudodifferential operators $\widehat{T}_1$, $\widehat{T}_2$ of disjoint compact microsupport, such that their principal symbol is 1 in a neighborhood $\overline{x}$, $\overline{y}$ respectively. Notice that

$$\widehat{T}_1\left[\Pi, \widehat{Q}\right]\widehat{T}_2 = \left[\widehat{T}_1\Pi\widehat{T}_2, \widehat{Q}\right] + \widehat{T}_1\Pi\left[\widehat{Q}, \widehat{T}_2\right] + \left[\widehat{Q}, \widehat{T}_1\right]\Pi\widehat{T}_2. \tag{5.12}$$

Near $(\overline{x}, \overline{y})$, the symbol of $\widehat{T}_1\left[\Pi, \widehat{Q}\right]\widehat{T}_2$ and $\left[\Pi, \widehat{Q}\right]$ coincide. Consider, on the other hand, the first term on the right-hand side of equation (5.12). First, by the assumption on the microsupports of T_1 and T_2, the operator $\widehat{T}_1\Pi\widehat{T}_2$ does not have wave-front set along the diagonal, and therefore it is in $\text{sc-}I^{-1/2}\left(M \times M; \mathcal{F}\partial X_c\right)$. We can then apply Proposition 6.3 below to compute the symbol of the commutator $\left[\widehat{T}_1\Pi\widehat{T}_2, \widehat{Q}\right]$. Near $(\overline{x}_0, \overline{y}_0)$ the symbol $\widehat{T}_1\Pi\widehat{T}_2$ is equal to the symbol of Π, and clearly the (diagonal) Lie derivative of this symbol with respect to the Hamilton flow of Q is zero. Therefore the symbol of this commutator (as an operator of order $-3/2$) is zero.

The principal symbols of the last two terms in (5.12) also vanish because the principal symbols of T_1, T_2 are constant near $\overline{x}_0, \overline{y}_0$, respectively. Therefore, the $(-3/2)$ principal symbol of $\left[\Pi, \widehat{Q}\right]$ vanishes off the diagonal, and therefore everywhere on $\mathcal{F}\partial X_c$ by continuity. This concludes the proof. □

6. On some propagators $e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}$

6.1. **The classical counterpart.** We begin by considering classical hamiltonians $Q : T^*M \to \mathbb{R}$ with the property that their Hamilton field is tangent to ∂X_c, that is, $\Xi_Q(\overline{x}) \in T_{\overline{x}}\partial X_c$ for all $\overline{x} \in \partial X_c$. It is easy to see that this occurs if and only if the Poisson bracket satisfies $\{P, Q\}_{|X_c} = 0$, and a as consequence, Q is constant on the fibers of $\partial X_c \to S$. The Hamilton flow of such a Q preserves the region X_c, that is, it defines a classical flow in the symplectic manifold with boundary X_c. The restriction of Q to ∂X_c descends to a smooth function $Q_S : S \to \mathbb{R}$, which in turn defines a Hamilton flow on S. However, the following diagram (where Φ_t^Q denotes the Hamilton flow of Q, etc.) does *not* commute in general:

$$\begin{tikzcd}
\partial X_c \arrow[r, "\Phi_t^Q|_{\partial X_c}"] \arrow[d] & \partial X_c \arrow[d] \\
S \arrow[r, "\Phi_t^{Q_S}"] & S
\end{tikzcd} \tag{6.1}$$

Lemma 6.1. *Assume that Ξ_Q is tangent to ∂X_c, and let P be a defining function of X_c, as above. Then the diagram (6.1) commutes for all t if and only if the Poisson bracket of P and Q vanishes to second order at the boundary ∂X_c, by which we mean that there exists a smooth function F such that*

$$\{P, Q\} = P^2 F.$$

Proof. Since $\Xi_Q(\overline{x}) \in T_{\overline{x}}\partial X_c$ for all $\overline{x} \in \partial X_c = P^{-1}(0)$, then $\{Q, P\}(\overline{x}) = dP_{\overline{x}}(\Xi_Q) = 0$ for all $\overline{x} \in \partial X_c$. Therefore we can write $\{P, Q\} = F_0 P$ for some smooth function F_0, and

$$[\Xi_P, \Xi_Q] = \Xi_{\{P,Q\}} = F_0 \Xi_P + P \Xi_{F_0}$$

will vanish on ∂X_c if and only if F_0 itself vanishes on ∂X_c. As a result,

$$\Phi_s^P \circ \Phi_t^Q(\overline{x}) = \Phi_t^Q \circ \Phi_s^P(\overline{x})$$

$\forall \overline{x} \in \partial X_c$ if and only if $\{P, Q\}$ vanishes to second order on ∂X_c. □

We now discuss the relation of the above considerations with Lerman's symplectic cut construction. The "cut" space is

$$Y = X_c / \sim,$$

where the equivalence relation on X_c is: $\overline{x} \sim \overline{y}$ iff $\overline{x}$ and $\overline{y}$ are on the boundary ∂X_c and in fact on the same leaf of $\partial X_c \to S$. There is an obvious inclusion $S \hookrightarrow Y$ and it is clear that, as sets,

$$Y = \mathrm{Int}\ (X_c) \coprod S \tag{6.2}$$

(disjoint union). Let us give Y the quotient topology. Then a function Q satisfying $\{Q, P\}|_{\partial X_c} = 0$ induces a continuous function $Q_Y : Y \to \mathbb{R}$, which is smooth when restricted to each of the pieces in (6.2). If the Poisson bracket vanishes to second order, then, by the previous lemma, one has a commutative diagram

$$\begin{tikzcd} X_c \arrow[r, "\Phi_t^Q|_{X_c}"] \arrow[d] & X_c \arrow[d] \\ Y \arrow[r, "\Phi_t^{Q_Y}"] & Y \end{tikzcd} \tag{6.3}$$

where Φ^{Q_Y} is a Hamilton flow defined piece-wise by restricting Q to the pieces in (6.2).

In Lerman's construction the topological space Y acquires the structure of a symplectic manifold of which S is a symplectic submanifold. However, in general, Q_Y *is not smooth* with respect to Lerman's structure; for this it is necessary that $\{Q, P\}|_{\partial X_c} = 0$ to infinite order, as implied by the following lemma:

Lemma 6.2. *If $\{P, Q\}|_{\partial X_c} = 0$ to order k, meaning that there exists a smooth function F such that $\{P, Q\} = P^k F$, then $Q_Y \in C^{k-1}(Y)$.*

Proof. Take the model case $M = \mathbb{R}^{n-1} \times S^1$ with coordinates (x, θ), T^*M with canonical coordinates $(x, \theta\,;\, p, \tau)$, and $X_c = \{(x, \theta\,;\, p, \tau) \big| \tau \geq 0\}$. Then the symplectic cut is the manifold $Y \cong \mathbb{R}^{2(n-1)}_{(x,p)} \times \mathbb{C}_z$ (where the symplectic form on $\mathbb{C}$ is $(\frac{1}{\sqrt{2}i} dz \wedge d\bar{z})$), and the projection $X_c \to Y$ is

$$(x, \theta; p, \tau) \mapsto (x, p; \sqrt{\tau} e^{i\theta}).$$

Any function $Q(x, \theta; p, \tau)$ whose Hamilton flow preserves X_c descends to the continuous function Q_Y on Y

$$Q_Y(x, p, z) = Q(x, \arg z\,;\, p, |z|^2)$$

for $z \neq 0$ and $Q_Y(x, p, 0) = Q(x, \theta\,;\, p, 0)$ for any value of θ. (The continuity of Q_Y will be seen in what follows.) If $\{P, Q\}$ vanishes to order k in ∂X_c, then

$$\{P, Q\} = \frac{\partial Q}{\partial \theta} = \tau^k\, F(x, \theta\,;\, p, \tau)$$

for some smooth function F. This implies that there exist smooth functions $G_1(x, \theta\,;\, p, \tau)$, $G_2(x\,; p, \tau)$ such that

$$Q = \tau^k\, G_1(x, \theta\,;\, p, \tau) + G_2(x\,;\, p, \tau).$$

It follows that

$$Q_Y(x, p, z) = \begin{cases} |z|^{2k}\, G_1(x, \arg z\,;\, p, |z|^2) + G_2(x\,;\, p, |z|^2) & \text{if } z \neq 0 \\ \\ G_2(x\,;\, p, 0) & \text{if } z = 0. \end{cases}$$

Define $\tilde{Q}_Y(x', \xi', z) = Q_Y(x', \xi', z) - G_2(x', |z|^2, \xi')$. Then Q_Y and $\tilde{Q}_Y$ differ by a smooth function, and therefore it suffices to show that $\tilde{Q}_Y(x', \xi', z)$ is C^k. For this, we'll show that a function of the form

$$\tilde{G}^k(x, p, z) = \begin{cases} |z|^{2k}\, G(x, \arg z\,;\, p, |z|^2) & \text{if } z \neq 0 \\ \\ 0 & \text{if } z = 0 \end{cases}$$

is a C^k function for any smooth function $G(x, \theta\,;\, p, \tau)$. The function $\tilde{G}^k$ is smooth in the region $z \neq 0$, so we only need to show the existence and continuity of partial derivatives at $z = 0$. We will prove the statement by induction. Write $z = (u_1, u_2)$. For $k = 1$,

$$\frac{\partial \tilde{G}^1(x, p, z)}{\partial u_1}\Big|_{z=0} = \lim_{u_1 \to 0} \frac{u_1^2 G(x, 0\,;\, p, u_1^2)}{u_1} = 0,$$

and for $z \neq 0$

$$\frac{\partial \tilde{G}^1}{\partial u_1} = 2u_1 G(x, \arg z\,;\, p, |z|^2) - u_2 \frac{\partial G}{\partial \theta}(x, \arg z\,;\, p, |z|^2) + 2u_1 |z|^2 \frac{\partial G}{\partial \tau}(x, \arg z\,;\, p, |z|^2),$$

which converges to zero as $z \to 0$. The partial derivative with respect to u_2 is similar, showing that $\tilde{G}^1 \in C^1$.

Assume that the statement is valid for $k-1>0$. Notice that

$$\frac{\partial \tilde{G}^k}{\partial u_1} = 2ku_1\tilde{G}^{k-1} - u_2 \widetilde{\left(\frac{\partial G}{\partial \theta}\right)}^{k-1} + 2u_1 \widetilde{\left(\frac{\partial G}{\partial \tau}\right)}^{k}, \text{ and}$$

$$\frac{\partial \tilde{G}^k}{\partial u_2} = 2ku_2\tilde{G}^{k-1} + u_1 \widetilde{\left(\frac{\partial G}{\partial \theta}\right)}^{k-1} + 2u_2 \widetilde{\left(\frac{\partial G}{\partial \tau}\right)}^{k}.$$

Each of the terms on the right-hand side of each of the equalities above are at least C^{k-1}, finishing the proof.

□

6.2. **A Symbolic Description of The Propagator $e^{-it\ h^{-1}\Pi\widehat{Q}\Pi}$.** Throughout this section Q will denote a smooth function such that the Poisson bracket $\{P,Q\}$ vanishes to second order at ∂X_c (c.f. Lemma 6.1). As we saw in the previous section we then obtain a classical flow $\Phi_t^Q|_{X_c}$ that descends to a continuous flow on the cut space Y. In this section we analyze the quantum mechanical propagator $e^{-it\ h^{-1}\Pi\widehat{Q}\Pi}$, where $\widehat{Q}$ is a semiclassical pseudodifferential operator on M with symbol Q and whose subprincipal symbol satisfies

$$\text{Sub } \widehat{Q}\Big|_{X_c} = 0.$$

Before we state the main result, let us start with a proposition:

Proposition 6.3. *For each semiclassical Fourier integral operator $\widetilde{V}(\hbar)$ in sc-$I^{-1/2}(M \times M; \mathcal{F}\partial X_c)$, the commutator $\left[\widehat{Q}(\hbar), \widetilde{V}(\hbar)\right]$ is in the class sc-$I^{-3/2}(M \times M; \mathcal{F}\partial X_c)$. Its principal symbol is*

$$\sigma_{[\widehat{Q},\widetilde{V}]}(\bar{x},\bar{y}) = \frac{\hbar}{i}\mathfrak{L}_{\Xi_Q} V(\bar{x},\bar{y}),$$

where V is the principal symbol of $\widetilde{V}$, and $\mathfrak{L}_{\Xi_Q}$ is the Lie derivative obtained by letting the Hamilton flow of Q act diagonally on $\mathcal{F}\partial X_c$.

Proof. Write the Schwartz kernel of $\widehat{Q}$ in the model case as

$$\widehat{Q}(y,x) = \frac{1}{(2\pi\hbar)^n}\int e^{i(y-x)p/h} q(y,p,\hbar)dp,$$

where

$$q(y,p,\hbar) \sim q_0(y,p) + \hbar q_1(y,p) + \ldots,$$

and the Schwartz kernel of $\widetilde{V} \in$ sc-$I^{-1/2}(M \times M\ ;\ \mathcal{F}\partial X_c)$ as

$$\widetilde{V}(z,y) = \frac{1}{(2\pi\hbar)^n}\int e^{i(z'-y')\omega/\hbar} v(z,y,\omega,\hbar)d\omega,$$

where

$$v(z,y,\omega,\hbar) \sim v_0(z,y,\omega) + \hbar v_1(z,y,\omega) + \dots$$

Then the Schwartz kernel of $\widetilde{V} \circ \widehat{Q}$ is

$$\widetilde{V} \circ \widehat{Q}(z,x) = \frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}(z'-x')\omega} \left[\frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(y-x)p+(x'-y')\omega]} v(z,y,\omega,\hbar) q(y,p,\hbar) dy dp \right] d\omega$$

Applying the stationary phase method to the integral in brackets, one gets

(6.4)

$$\begin{aligned} &\frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(y-x)p+(x'-y')\omega]} v(z,y,\omega,\hbar) q(y,p,\hbar) dy dp \\ &\sim v_0(z,x,\omega) q_0(x,(0,\omega)) - \frac{\hbar}{i} \frac{\partial v_0(z,x,\omega)}{\partial x} \frac{\partial q_0(x,p)}{\partial p}\Big|_{p=(0,\omega)} - \frac{\hbar}{i} v_0(z,x,\omega) \frac{\partial q_0(x,p)}{\partial x \partial p}\Big|_{p=(0,\omega)} \\ &+ \hbar v_1(z,x,\omega) q_0(x,(0,\omega)) + \hbar v_0(z,x,\omega) q_1(x,(0,\omega)). \end{aligned}$$

The Schwartz kernel of $\widehat{Q} \circ \widetilde{V}$ can be written as

$$\widehat{Q} \circ \widetilde{V}(z,x) = \frac{1}{(2\pi\hbar)^n} \int e^{i(z'-x')\omega/\hbar} \left[\frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(z-y)p+(y'-z')\omega]} q(z,p,\hbar) v(y,x,\omega,\hbar) dp dy \right] d\omega$$

Applying stationary phase to the amplitude above, one gets

$$\begin{aligned} &\frac{1}{(2\pi\hbar)^n} \int e^{i\hbar^{-1}[(z-y)p+(y'-z')\omega]} q(z,p,\hbar) v(y,x,\omega,\hbar) dp dy \\ &\sim q_0(z,(0,\omega)) v_0(z,x,\omega) + \frac{\hbar}{i} \frac{\partial q_0(z,p)}{\partial p}\Big|_{p=(0,\omega)} \frac{\partial v_0(z,x,\omega)}{\partial z} \\ &+ \hbar q_1(z,(0,\omega)) v_0(z,x,\omega) + \hbar q_0(z,(0,\omega)) v_1(z,x,\omega) \end{aligned} \tag{6.5}$$

Therefore, the commutator $[\widehat{Q}, \widetilde{V}]$ has as leading amplitude

$$v_0(z,x,\omega) \left(q_0(z,(0,\omega) - q_0(x,(0,\omega)) \right).$$

This vanishes at $x' = z'$, which corresponds to the flow-out. Therefore $[\widehat{Q}, \widetilde{V}] \in$ sc-$I^{-3/2}(M \times M \ ; \ \mathcal{F}\partial X_c)$. In order to compute the principal symbol there, we notice that

$$q_0(z,(0,\omega)) - q_0(x,(0,\omega)) = (z'-x') \cdot d(z,x,\omega),$$

where $d = (d_2, \dots, d_n)$ is a vector-valued function such that

$$d(z,x,\omega)\Big|_{x'=z'} = \nabla_{x'} q_0(x,(0,\omega))\Big|_{x'=z'}.$$

Therefore

$$\frac{1}{(2\pi\hbar)^n}\int e^{i(z'-x')\omega/\hbar}\left(q_0(z,(0,\omega)-q_0(x,(0,\omega))\right)v_0(z,x,\omega)d\omega$$

$$= \frac{-\hbar}{i(2\pi\hbar)^n}\int e^{i(z'-x')\omega/\hbar}\sum_{j=1}^{n-1}\frac{\partial}{\partial\omega_j}\left(d_j(z,x,\omega)v_0(z,x,\omega)\right)d\omega. \tag{6.6}$$

The principal symbol of $[\widehat{Q},\widetilde{V}]$ in sc-$I^{-3/2}(M\times M\ ;\ \mathcal{F}\partial X_c)$ can be then computed taking all the contributions from equations (6.4), (6.5) and (6.6). Since $q_0(x,p)$ is constant along the orbits, then

$$\frac{\partial^2 q_0(x,p)}{\partial x_j\partial p_j}\Big|_{p=(0,\omega)} = \frac{\partial^2 q_0(z,p)}{\partial x_j\partial p_j}\Big|_{\substack{x'=z'\\ p=(0,\omega)}},\ \text{for any } j\geq 2.$$

Since $\{Q,P\}$ vanishes at second order on ∂X_c, then $\frac{\partial^2 q_0(x,p)}{\partial x_1\partial x_1}\Big|_{p_1=0} = 0$. Assuming all these conditions, and the fact that the subprincipal symbol

$$\text{Sub }\widehat{Q}(x,p) = q_1(x,p) - \frac{1}{2i}\sum_{j=1}^{n}\frac{\partial^2}{\partial x_j\partial p_1}\Big|_{p=(0,\omega)}$$

vanishes on ∂X_c, the principal symbol of the commutator reduces to

$$\frac{\partial q_0(z,p)}{\partial p}\Big|_{\substack{x'=z'\\ p=(0,\omega)}}\frac{\partial v_0(z,x,\omega)}{\partial z}\Big|_{x'=z'} + \frac{\partial q_0(x,p)}{\partial p}\Big|_{\substack{x'=z'\\ p=(0,\omega)}}\frac{\partial v_0(z,x,\omega)}{\partial x}\Big|_{x'=z'}$$
$$-\sum_{j=2}^{n}\frac{\partial q_0(x,p)}{\partial x_j}\Big|_{\substack{x'=z'\\ p=(0,\omega)}}\frac{\partial v_0(z,x,\omega)}{\partial \omega}\Big|_{x'=z'}.$$

One can check that this is the Lie derivative of V with respect to the Hamiltonian field Ξ_Q in the sense in the statement of the lemma. □

Notice that there is a way to define classes $J^{\ell,m}$ for a general pair of admissible Lagrangian submanifolds that intersect cleanly (see [22, 11]), and not only for the diagonal and the flow-out of ∂X_c. The main result of this section is the following:

Theorem 6.4. *Suppose $\widehat{Q}$ is a zeroth-order semiclassical pseudodifferential operator satisfying the conditions of Lemma 6.1. Assume sub$\widehat{Q}(\hbar) = 0$. Then*

$$\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi} \in J^{-1/2,1/2}\left(M\times M;\Delta(t),\mathcal{F}\partial X_c(t)\right),$$

where

$$\Delta(t) = \left\{(\overline{x},\overline{y})\,\middle|\,\overline{x},\overline{y}\in T^*(M\times M), \overline{x} = \Phi_t^Q(\overline{y})\right\} \tag{6.7}$$

$$\mathcal{F}\partial X_c(t)=\left\{(\overline{x},\overline{y})\,\middle|\,\overline{x},\overline{y}\in\partial X_c,\ \exists s\in\mathbb{R}\ \textit{such that}\ \overline{x}=\Phi^P_s\Phi^Q_t(\overline{y})\right\}. \tag{6.8}$$

Remark 6.5. In this statement t is a parameter, but we could also consider t as a variable (in which case the kernel of the operator would be a family of functions on $\mathbb{R}\times M\times M$). Also, the symbols of $\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}$ can easily be computed.

Proof. Let us define the following operator:

$$W(t):=\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}\,e^{it\hbar^{-1}\widehat{Q}}. \tag{6.9}$$

We first prove the following:

Lemma 6.6. *$W(t)\in J^{-1/2,1/2}\left(M\times M;\Delta,\mathcal{F}\partial X_c\right)$, and the principal symbol in the diagonal is $\sigma_0=\chi_{X_c}$.*

Proof. Let us define $D_t=\frac{1}{i}\frac{\partial}{\partial t}$. $W(t)$ satisfies the following equation

$$\begin{cases}\hbar D_tW(t)+\left[\widehat{Q},W(t)\right]+\left[\Pi,\widehat{Q}\right]W(t)=0\\[2ex] W\Big|_{t=0}=\Pi\end{cases} \tag{6.10}$$

Similarly to the proof of Lemma 5.8, using the symbol calculus we will construct a sequence of approximate solutions of equation (6.10). This construction makes the right-hand side of order $O(\hbar^\infty)$, and an application of Duhamel's principle concludes the proof.

As a first approximation we take $\widetilde{W_0}=\Pi$. This is a sensible choice since

$$\hbar^2\widetilde{S}_2:=\hbar D_t\widetilde{W_0}+\left[\widehat{Q},\widetilde{W_0}\right]+\left[\Pi,\widehat{Q}\right]\widetilde{W_0}=-\left[\Pi,\widehat{Q}\right](I-\Pi)\in\text{sc-}I^{-5/2}\left(M\times M;\mathcal{F}\partial X_c\right),$$

by Proposition 5.11.

We now modify $\widetilde{W_0}$ by elements in $I(M\times M;\mathcal{F}\partial X_c)$ to lower the order of the remainder. It is easy to see that the symbol of the correction term is the solution to the problem

$$\begin{cases}\frac{\partial V_1(\bar{x},\bar{y},t)}{\partial t}+\mathfrak{L}_{\Xi_Q}V_1(\bar{x},\bar{y},t)=-iS_2,\\[2ex] V_1\Big|_{t=0}=0\end{cases}$$

where S_1 is the principal symbol of $\widetilde{S}_1\in\text{sc-}I^{-1/2}(M\times M;\mathcal{F}\partial X_c)$. Let $\widetilde{V}_1\in\text{sc-}I^{-1/2}(M\times M;\mathcal{F}\partial X_c)$ be an operator with this as symbol, and let $\widetilde{W_1}=\widetilde{W_0}+\hbar\widetilde{V}_1$.

Since $\left[\Pi,\widehat{Q}\right]\in\text{sc-}I^{-5/2}(M\times M;\mathcal{F}\partial X_c)$ by Proposition 5.11, we get

$$\hbar^3\widetilde{S}_3:=\hbar D_t\widetilde{W_1}+\left[\widehat{Q},\widetilde{W_1}\right]+\left[\Pi,\widehat{Q}\right]\widetilde{W_1}\in I^{-7/2}(M\times M;\mathcal{F}\partial X_c)$$

Proceeding inductively in this fashion, we obtain an infinite sequence $\{\widetilde{V}_j\}$ such that for all J

$$\hbar D_t \left(\widetilde{W_0} + \sum_{j=1}^{J} \hbar^j \widetilde{V}_j \right) + \left[\widehat{Q}, \widetilde{W_0} + \sum_{j=1}^{J} \hbar^j \widetilde{V}_j \right] + \left[\Pi, \widehat{Q} \right] \left(\widetilde{W_0} + \sum_{j=1}^{J} \hbar^j \widetilde{V}_j \right)$$

$$= \hbar^{J+2} \widetilde{S}_{J+2} \in I^{-5/2-J}(M \times M; \mathcal{F}\partial X_c).$$

Next we take an operator $\widetilde{V} \in \text{sc-}I^{-1/2}\left(M \times M; \mathcal{F}\partial X_c\right)$ such that $\widetilde{V} \sim \sum_{j=1}^{\infty} \hbar^j \widetilde{V}_j$, and define $\widetilde{W} = \widetilde{W_0} + \widetilde{V}$. $\square$

Going back to the proof of the theorem, notice that

$$\Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi} = W(t) e^{-itN\widehat{Q}}.$$

The Lagrangian $\Delta(t)$ intersects Δ and $\mathcal{F}\partial X_c$ transversally. Using a variation of the Proposition 4.1 in [11], we conclude that composing elements in $J^{-1/2,1/2}(M \times M; \Delta, \mathcal{F}\partial X_c)$ with $e^{-it\hbar^{-1}\widehat{Q}}$ gives elements in $J^{-1/2,1/2}(M \times M; \Delta(t), \mathcal{F}\partial X_c(t))$. $\square$

6.3. **An Egorov-type Theorem.** We can easily prove the following corollary.

Corollary 6.7. *Let $\widehat{Q}$ be a zeroth order semiclassical pseudodifferential operator satisfying the conditions of Lemma 6.1, and the Levi condition. Then for any zeroth order semiclassical pseudodifferential operator $\widehat{A}(\hbar)$, we have*

$$\widetilde{B}(t) := e^{it\hbar^{-1}\Pi\widehat{Q}\Pi} \Pi \widehat{A} \Pi e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi} \in J^{-1/2,1/2}\left(M \times M, \Delta, \mathcal{F}\partial X_c\right),$$

with the following principal symbols:

$$\sigma_0(\widetilde{B}(t))(\overline{x}, \overline{x}) = \chi_{X_c}(\overline{x}) \, \left(a \circ \Phi_t^Q(\overline{x}) \right) \quad \textit{for } (\overline{x}, \overline{x}) \in \Delta \setminus \Sigma, \textit{and}$$

$$\sigma_1(\widetilde{B}(t))\Big|_{F_s} = \Pi_{F_s} M_{(a \circ \Phi_t^Q)_{|F_s}} \Pi_{F_s},$$

where a is the principal symbol of $\widehat{A}$, Φ_t^Q the Hamilton flow Q, F_s is an orbit in ∂X_c, and Π_{F_s} is the Szegö projector of F_s.

Proof. Let us consider $W(t)$ as in equation (6.9). Notice that

$$\widetilde{B}(t) = e^{it\hbar^{-1}\Pi\widehat{Q}\Pi} \Pi \widehat{A} \Pi e^{-itN\Pi\widehat{Q}\Pi} = W(-t) e^{it\widehat{Q}} \widehat{A} e^{-it\widehat{Q}} W(-t)^*,$$

which proves it belongs to the algebra. Since $e^{it\hbar^{-1}\widehat{Q}} \widehat{A} e^{-it\hbar^{-1}\widehat{Q}}$ is a semiclassical pseudodifferential operator with symbol $a \circ \Phi_t^Q$, the symbol on the diagonal can be trivially obtained. To compute the symbol on the flow-out, we note that the principal symbols of W are exactly those of Π and we use Proposition 2.7:

$$\sigma_1(\widetilde{B}(t))(\overline{y} = \Phi_s^P(\overline{x}), \overline{x}) =$$

$$= \frac{1}{\sqrt{2\pi}} \int \sigma_1\left(W(-t)\right)(\overline{y}, \Phi^P_{\tilde{s}}(\overline{x}))\sigma_1\left(e^{it\widehat{Q}}\widehat{A}e^{-it\widehat{Q}}W(-t)^*\right)(\Phi^P_{\tilde{s}}(\overline{x}),\overline{x})d\tilde{s}$$

$$= \frac{1}{\sqrt{2\pi}}\int \frac{1}{\sqrt{2\pi}}\frac{1}{1-e^{i(s-\tilde{s})}} a(\Phi^Q_t\Phi^P_{\tilde{s}}(\overline{x}))\frac{1}{\sqrt{2\pi}}\frac{1}{1-e^{i\tilde{s}}}d\tilde{s}.$$

As a pseudodifferential operator on the fibers, this is $\Pi_{F_s} M_{(a\circ\Phi^Q_t)_{|F_s}}\Pi_{F_s}$. □

7. A NUMERICAL STUDY OF PROPAGATION OF COHERENT STATES

Let $\widehat{Q}$ be a zeroth order semiclassical pseudodifferential operator with symbol Q. It is well-known that, if $\psi_{(x_0,p_0)}$ is a coherent state with center at (x_0,p_0), then $e^{-it\hbar^{-1}\widehat{Q}}(\psi_{(x_0,p_0)})$ is a coherent state (appropriately "squeezed") with center at $\Phi^Q_t(x_0,p_0)$, where Φ^Q is the Hamilton flow of Q. If the flow Φ^Q preserves X_c and the center (x_0,p_0) is in the interior of X_c, then the same conclusion holds for the propagation $e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}(\psi_{(x_0,p_0)})$ of the coherent state by $\Pi\widehat{Q}\Pi$, as the trajectory of the center will remain away from the boundary ∂X_c and everything is as if we were in the boundaryless case.

In this section we present results of a numerical calculation of $e^{-it\hbar^{-1}\Pi\widehat{Q}\Pi}(\psi_{(x_0,p_0)})$ in an example where the Hamilton flow of Q does not preserve X_c, that is, trajectories of Φ^Q cross the boundary ∂X_c.

We consider the Harmonic oscillator $\widehat{P} = \frac{1}{2}(x^2 - \hbar^2\partial_x^2)$ in $\mathbb{R}^1$, and the corresponding projector Π onto the span of its eigenfunctions of with eigenvalues less than or equal to one. We take $Q = x^2 - p^2$, and $\widehat{Q}$ the obvious quantization of Q. Figure 1 shows some energy levels of Q. Notice that the energy levels cross the boundary of $X_c = \left\{x^2 + p^2 \leq 2\right\}$.



FIGURE 1. Energy levels of Q and the boundary $\partial X_c = \{P = 1\}$.

We now take a coherent state centered inside the interior of X_c, and numerically compute its propagation under $\Pi\widehat{Q}\Pi$. We do the calculation in Bargmann space, for symplicity.

We recall that the Bargmann space is defined as the Hilbert space

$$\mathcal{B}=\left\{f:\mathbb{C}\to\mathbb{C}: f \text{ entire and } \int |f(z)|^2\, e^{-\frac{|z|^2}{\hbar}} dm_z<\infty\right\},$$

where $dm_z = dxdp$ and $z=\frac{x-ip}{\sqrt{2}}$, with the Hermitian inner product

$$\langle f,g\rangle=\int f(z)\overline{g(z)}e^{-\frac{z\overline{z}}{\hbar}}dm_z.$$

The Harmonic oscillator in Bargmann space is given by

$$\widehat{P}=\hbar z\frac{\partial}{\partial z}+\frac{\hbar}{2},$$

with principal symbol $P(z,\overline{z})=z\overline{z}$ and eigenbasis

$$\left\{b_n=\frac{z^n}{\sqrt{\hbar^n n!}}\right\}_{n\in\mathbb{N}\cup\{0\}},\ \widehat{P}b_n=\hbar\left(n+\frac{1}{2}\right)b_n=\lambda_n b_n.$$

The quantization of $Q=x^2-p^2$ in Bargmann space is the operator $\widehat{Q}=\hbar^2\frac{\partial^2}{\partial z^2}+z^2$. Applying $\widehat{Q}$ to the eigenbasis, we get

$$\widehat{Q}(b_n)=\hbar\sqrt{n(n-1)}b_{n-2}+\hbar\sqrt{(n+1)(n+2)}b_{n+2},$$

which gives a a "generalized" Toeplitz matrix for $\Pi_N\widehat{Q}\Pi_N$ for each positive integer N, where $\hbar=\frac{1}{N}$. The (normalized) coherent state in Bargmann space, with center at w, is given by the simple formula

$$\Psi_w(z)=e^{\frac{z\overline{w}}{\hbar}}e^{-\frac{w\overline{w}}{2\hbar}}.$$

We apply the propagator $e^{-it\hbar^{-1}\Pi_N\widehat{Q}\Pi_N}$ to the projected coherent state

$$\Psi_w(z,N):=\Pi_N\Psi_w(z)=\sum_{n=0}^{N}\frac{\overline{w}^n}{\sqrt{n!\hbar^n}}e^{-\frac{\overline{w}w}{2\hbar}}b_n.$$

In Bargmann space, we measure the concentration in phase space of any semiclassical family ψ by taking the absolute value of the family times the square root of the Bargmann weight, namely, by forming the Husimi density:

$$|\psi|_H(z):=|\psi(z)|\,e^{-z\overline{z}/2\hbar}.$$

We took as initial data a projected coherent state with center at $w=-0.25-0.6i$, which corresponds to $(x,p)=\sqrt{2}\,(-0.25,0.6)$. Figure 2 consists of contour plots of the Husimi density in the z variable of the initial projected coherent state, its propagation at $t=0.25$ (approximately when the center of the coherent state hits the boundary), and at time $t=0.5$. We observe that after the time of collision of the center with the boundary, the coherent state splits into two localized states with centers on inward trajectories and with the same classical energy. Here we took $N=100$. The splitting happens immediately after the center collides with the



Figure 2. A coherent state is propagated by $e^{-itN\Pi_N\hat{Q}\Pi_N}$ ($Q = x^2 - p^2$) at time $t = 0$ (left), $t = .25$ (middle) and $t = 0.5$ (right). The contour plots of the Husimi densities at each time are shown.

boundary; thus one can speak of infinite-propagation speed along the boundary. Note that the evolution is time-reversible, so that in some cases the opposite phenomenon will occur, namely, two localized states with same classical energy will hit the boundary at the same time and combine into one.

References

[1] José L. Antoniano and Gunther A. Uhlmann. A functional calculus for a class of pseudodifferential operators with singular symbols. In *Pseudodifferential operators and applications (Notre Dame, Ind., 1984)*, volume 43 of *Proc. Sympos. Pure Math.*, pages 5–16. Amer. Math. Soc., Providence, RI, 1985.

[2] Martin Bojowald and Thomas Strobl. Group theoretical quantization and the example of a phase space $S^1 \times \mathbf{R}^+$. *J. Math. Phys.*, 41(5):2537–2567, 2000.

[3] Martin Bojowald and Thomas Strobl. Symplectic and projection quantization for nonholonomic constraints. *Internat. J. Modern Phys. D*, 12(4):713–725, 2003.

[4] Sandrine Dozias. Clustering for the spectrum of h-pseudodifferential operators with periodic flow on an energy surface. *J. Funct. Anal.*, 145(2):296–311, 1997.

[5] Dimitri Gioev. Lower order terms in Szegö type limit theorems on Zoll manifolds. *Comm. Partial Differential Equations*, 28(9-10):1739–1785, 2003.

[6] Ulf Grenander and Gabor Szegö. *Toeplitz forms and their applications*. California Monographs in Mathematical Sciences. University of California Press, Berkeley, 1958.

[7] V. Guillemin and E. Lerman. Melrose-Uhlmann projectors, the metaplectic representation and symplectic cuts. *J. Differential Geom.*, 61(3):365–396, 2002.

[8] V. Guillemin and K. Okikiolu. Spectral asymptotics of Toeplitz operators on Zoll manifolds. *J. Funct. Anal.*, 146(2):496–516, 1997.

[9] V. Guillemin and K. Okikiolu. Subprincipal terms in Szegö estimates. *Math. Res. Lett.*, 4(1):173–179, 1997.

[10] V. Guillemin and S. Sternberg. *Semi-classical analysis*. Preprint, MIT, 2010.

[11] V. Guillemin and G. Uhlmann. Oscillatory integrals with singular symbols. *Duke Math. J.*, 48(1):251–267, 1981.

[12] Victor Guillemin. Some classical theorems in spectral theory revisited. In *Seminar on Singularities of Solutions of Linear Partial Differential Equations (Inst. Adv. Study, Princeton,*

N.J., 1977/78), volume 91 of *Ann. of Math. Stud.*, pages 219–259. Princeton Univ. Press, Princeton, N.J., 1979.

[13] Victor Guillemin. Gauged Lagrangian distributions II. *preprint (MIT)*, 1994.

[14] Victor Guillemin and Shlomo Sternberg. *Geometric asymptotics*. American Mathematical Society, Providence, R.I., 1977. Mathematical Surveys, No. 14.

[15] Victor Guillemin and Shlomo Sternberg. Some problems in integral geometry and some related problems in microlocal analysis. *Amer. J. Math.*, 101(4):915–955, 1979.

[16] Andrew Hassell and András Vasy. Intersecting Legendrians and blow-ups. *Math. Res. Lett.*, 8(4):413–428, 2001.

[17] B. Helffer and D. Robert. Puits de potentiel généralisés et asymptotique semi-classique. *Ann. Inst. H. Poincaré Phys. Théor.*, 41(3):291–331, 1984.

[18] Lars Hörmander. *The analysis of linear partial differential operators. I*, volume 256 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1983. Distribution theory and Fourier analysis.

[19] M. S. Joshi. A symbolic construction of the forward fundamental solution of the wave operator. *Comm. Partial Differential Equations*, 23(7-8):1349–1417, 1998.

[20] A. Laptev, D. Robert, and Yu. Safarov. Remarks on the paper of V. Guillemin and K. Okikiolu: "Subprincipal terms in Szegö estimates" [Math. Res. Lett. **4** (1997), no. 1, 173–179; MR1432819 (97k:58169)]. *Math. Res. Lett.*, 5(1-2):57–61, 1998.

[21] A. Laptev and Yu. Safarov. Szegö o type limit theorems. *J. Funct. Anal.*, 138(2):544–559, 1996.

[22] R. B. Melrose and G. A. Uhlmann. Lagrangian intersection and the Cauchy problem. *Comm. Pure Appl. Math.*, 32(4):483–519, 1979.

[23] T. Paul and A. Uribe. The semi-classical trace formula and propagation of wave packets. *J. Funct. Anal.*, 132(1):192–249, 1995.

[24] Harold Widom. Toeplitz determinants with singular generating functions. *Amer. J. Math.*, 95:333–383, 1973.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MICHIGAN 48109-1043.
E-mail address: `gerahdez@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MICHIGAN 48109-1043.
E-mail address: `uribe@umich.edu`